

Uroteuthis chinensis**Mitre squid****Scientific classification**

Kingdom: [Animalia](#)
 Phylum: [Mollusca](#)
 Class: [Cephalopoda](#)
 Order: [Myopsida](#)
 Family: [Loliginidae](#)
 Genus: [Uroteuthis](#)

Binomial name

Uroteuthis chinensis
 (Gray, 1849)


Synonyms^[2]

- *Loligo chinensis* Gray, 1849
- *Loligo etheridgei* Berry, 1918
- *Loligo formosana* Sasaki, 1929

A. Environment/Ecology:

Demersal; non-migratory (Ref. [75930](#)); depth range 15 - 170 m (Ref. [275](#)), usually 40 - 150 m (Ref. [75930](#)). Tropical; 21°C - 29°C (Ref. [75934](#)); 34°N - 30°S, 99°E - 154°E (Ref. [275](#))

B. Distribution:



Note: Distribution range colours indicate degree of suitability of habitat which can be interpreted as probabilities of occurrence.

Indo-West Pacific.

C. Length at first maturity / Size / Weight / Age:

Maturity: L_m ?, range 6 - 8 cm Max length : 30.0 cm ML male/unsexed; (Ref. [275](#))

D. Short description

Uroteuthis is a genus of 14 species of common inshore squids of the Indo-West Pacific and is further subdivided into 3 subgenera. The members of the genus *Uroteuthis* are the only squids of the family Loliginidae that possess photophores (light-emitting organs) and all species in the genus have a pair of photophore organs on the ventral surface of their ink sac either side of their intestine.

Uroteuthis species range in size between 3 cm to 100 cm (mantle length). As with all other members of the family Loliginidae, they have a cornea that covers the lens of each eye, and have a gladius that extends the full length of the mantle and a gill that has a branchial canal.

E. Biology

Also caught by scoop nets and bamboo stake nets. Members of the class Cephalopoda are gonochoric. Male and female adults usually die shortly after spawning and brooding, respectively. Mating behavior: Males perform various displays to attract potential females for copulation. During copulation, male grasp the female and inserts the hectocotylus into the female's mantle cavity where fertilization usually occurs. Life cycle: Embryos hatch into planktonic stage and live for some time before they grow larger and take up a benthic existence as adults (Ref. [833](#)).

F. Life cycle and mating behavior

Members of the class Cephalopoda are gonochoric. Male and female adults usually die shortly after spawning and brooding, respectively. Mating behavior: Males perform various displays to attract potential females for copulation. During copulation, male grasp the female and inserts the hectocotylus into the female's mantle cavity where fertilization usually occurs. Life cycle: Embryos hatch into planktonic stage and live for some time before they grow larger and take up a benthic existence as adults.

G. Fisheries

The mitre squid is both targeted and taken as bycatch throughout its range, but is not recorded separately in landing statistics. Jereb et al. (2010) estimate that this species comprises half the squid catch in the South China Sea, while Arkhipkin et al. (2015) estimate that it accounts for up to 90% of the total Chinese loliginid catch with a maximum catch of 100,000 tonnes. It is the predominant squid species in the Taiwan Strait (Liao et al. 2010) and is seasonally very common in Taiwanese fish markets (C.C. Lu, personal communication). This species represents about 76-80% of loliginid squid landings from squid light luring fisheries in Thai Waters (Songjitsawat and Sookbuntoeng 1988) and is the most commonly caught species in the East Indian Ocean (Arkhipkin et al. 2015). There are three main fishery areas in the South China Sea, each with different fishing seasons. The southern part of Hainan Island is fished April to September. The southwest part of Beibu Gulf is fished April to January. The Taiwan shoal is fished April to September. There is some variation in catch per unit effort across years (Arkhipkin et al. 2015), but no real trend.

H. IUCN Red List Status

GEOGRAPHIC RANGE

- **Taxonomy**

| | |
|----------|-----------------------------|
| Kingdom: | Animalia |
| Phylum: | Mollusca |
| Class: | Cephalopoda |
| Order: | Myopsida |
| Family: | Loliginidae |
| Genus: | Uroteuthis |

- **Geographic Range**

NUMBER OF LOCATIONS

UPPER DEPTH LIMIT : 10 metres

LOWER DEPTH LIMIT : 170 metres

RANGE DESCRIPTION

The mitre squid is widely distributed in coastal waters throughout the Indo-west Pacific. It has a depth range of 10-170 m but is most commonly found in the 30-50 m depth bracket (Arkhipkin et al. 2015). It is reported from the southern tip of India and Sri Lanka and throughout the Bay of Bengal, extending southwards down the Malay Peninsula to

Indonesia. It has been recorded throughout the Indo-west Pacific as far east as Papua New Guinea, southwards to northern, eastern and western coasts of Australia, and northwards to the Philippines, the South China Sea, the Gulf of Thailand, and coastal waters of continental Asia as far north and east as Japan and southeast Russia (Chotiyaputta et al. 1992, Natewathana 1992, Jereb et al. 2010).

- **Population**

CURRENT POPULATION TREND : *Unknow*

- **Habitat and Ecology**

System : Marine

Habitat type : Marine Neritic

- **Biological resource use :**

Fishing & harvesting aquatic resource

- **Threats**

Fishing is a potential threat to this species.

- **Use and Trade**

The mitre squid is both targeted and taken as bycatch throughout its range, but is not recorded separately in landing statistics. Jereb et al. (2010) estimate that this species comprises half the squid catch in the South China Sea, while Arkhipkin et al. (2015) estimate that it accounts for up to 90% of the total Chinese loliginid catch with a maximum catch of 100,000 tonnes. It is the predominant squid species in the Taiwan Strait (Liao et al. 2010) and is seasonally very common in Taiwanese fish markets (C.C. Lu, personal communication). This species represents about 76-80% of loliginid squid landings from squid light luring fisheries in Thai Waters (Songjitsawat and Sookbuntoeng 1988) and is the most commonly caught species in the East Indian Ocean (Arkhipkin et al. 2015). There are three main fishery areas in the South China Sea, each with different fishing seasons. The southern part of Hainan Island is fished April to September. The southwest part of Beibu Gulf is fished April to January. The Taiwan shoal is fished April to September. There is some variation in catch per unit effort across years (Arkhipkin et al. 2015), but no real trend.

- **Conservation Action**

There is a fishing ban in spawning grounds during the breeding season (Arkhipkin et al. 2015); it is not known whether any further conservation measures are needed. Research is required on the life history of the mitre squid, its population status and trends, and the impacts of fishing throughout its range.

I. More Information:

1) Stocks

There is some variation in catch per unit effort across years (Arkhipkin et al. 2015), but no real trend. There is no information on the population status and trends of this species.

2) Ecology

Ecology of *Uroteuthis chinensis*

Substrate

| | |
|------------------|----------------------------|
| Substrate | Benthic: mobile; demersal; |
|------------------|----------------------------|

3) Diet

Feeding

| | | | | | | |
|-----------------------------------|---|------------------------|------|----------------------------|------|--|
| feeding type | mainly animals (troph. 2.8 and up) | | | | | |
| feeding type ref | Yunrong, Y., L. Yuyuan, Y. Shengyun, W. Guirong, T. Yajin, F. Qibin and L. Huosheng, 2013 | | | | | |
| feeding habit | hunting macrofauna (predator) | | | | | |
| feeding habit ref | Yunrong, Y., L. Yuyuan, Y. Shengyun, W. Guirong, T. Yajin, F. Qibin and L. Huosheng, 2013 | | | | | |
| trophic level(s) | | original sample | | unfished population | | Remark |
| estimation method | | Troph | s.e. | Troph | s.e. | |
| From diet composition | | | | | | |
| Ref. | | | | | | |
| From individual food items | | 3.87 | 0.47 | | | Trophic level estimated from a number of food items using a randomized resampling routine. |

4) Reproduction

Reproduction of *Uroteuthis chinensis*

| | |
|--|---|
| Main Ref. | Ruppert, E.E., R.S. Fox and R.D. Barnes, 2004 |
| Mode | dioecism |
| Batch Spawner | No |
| Description of life cycle and mating behavior | Members of the class Cephalopoda are gonochoric. Male and female adults usually die shortly after spawning and brooding, respectively. Mating behavior: Males perform various displays to |

| | |
|---|---|
| | attract potential females for copulation. During copulation, male grasp the female and inserts the hectocotylus into the female's mantle cavity where fertilization usually occurs. Life cycle: Embryos hatch into planktonic stage and live for some time before they grow larger and take up a benthic existence as adults. |
| Search for more references on reproduction | Scirus |

5) Maturity

| Maturity studies for <i>Uroteuthis chinensis</i> | | | | | | | |
|---|-------------|---|---------------|--------|------------------------|---------|----------------------|
| n = 2 | | | | | | | |
| Sort by <input checked="" type="radio"/> Lm <input type="radio"/> Country <input type="radio"/> Locality <input type="radio"/> tm | | | | | | | |
| Lm (cm) | Length (cm) | | Age range (y) | tm (y) | Sex of fish | Country | Locality |
| | 5.5 | - | 8.0 | - | female | China | Not specified, China |
| 14.1 ML | 4.6 | - | 12.8 | - | male | China | Not specified, China |

6) Spawning

| Spawning for <i>Uroteuthis chinensis</i> |
|---|
| The mitre squid is a large-sized species: males grow to 49 cm mantle length, females to 31 cm mantle length, although they are commonly smaller than this. Males and females reach maturity at around 16 cm and 14 cm mantle length respectively, and maturity appears to be related to size rather than age. The sex ratio is 1:1.5 males to females (Arkhipkin et al. 2015). They spawn throughout the year but seasonal peaks in spawning activity are found. In Thai waters, these occur in March-June and August-November (Chotiyaputta 1995). |

7) Spawning aggregation

(NA)

8) Fecundity

Fecundity ranges from 3,000 to 20,000 eggs (Arkhipkin et al. 2015).

9) Eggs

(NA)

10) Egg development

(NA)

11) Age/Size

(NA)

12) Growth

Growth parameters for *Uroteuthis chinensis*

Maximum Length 30cm ML

n = 3

Note that studies where Loo is very different (+/- 1/3) from Lmax are doubtful.

[Auximetric graph](#) [n = 1] $\phi = 3.46$ L inf = 42.0 cm ML K = 1.7 Median record no. 2 Ref. [94032](#)

| Loo (cm) | Length Type | K (1/y) | Temp° C | ϕ' | Country | Locality | Questionable | Captive |
|----------|-------------|-----------------------|---------|---------|----------|-------------------|--------------|---------|
| 40.90 | DL | 0.490 | | 2.91 | | Gulf of Thailand | No | No |
| 42.00 | ML | 1.650 | 29.50 | 3.46 | Thailand | Gulf of Thailand | Yes | No |
| 44.10 | ML | 1.730 | 29.40 | 3.53 | Thailand | Andaman Sea Coast | Yes | No |

13) Length-weight

Length-Weight Parameters for *Uroteuthis chinensis*[Length-weight \(a vs b\) graph](#)

[n=10]

Median Record No. 6

a = 0.2132 cm ML

b = 2.2148

Sort by



b



Country



Locality

| a | b | Doubtful? | Sex | Length (cm) | Length type | No. | Country | Locality |
|------------------------|-------|-----------|---------|-------------|-------------|-----|-----------|-------------------------------------|
| 1.1523 | 1.631 | No | female | | ML | 450 | Indonesia | Brondong |
| 0.9693 | 1.684 | No | mixed | | ML | 882 | Indonesia | Brondong |
| 0.7794 | 1.754 | No | male | | ML | 432 | Indonesia | Brondong |
| 0.5210 | 1.803 | No | male | 8.4 - 37.0 | ML | 169 | Indonesia | Bangka Regency |
| 0.3964 | 1.820 | Yes | male | 5.0 - 28.6 | ML | 91 | Thailand | Andaman Sea / 2005-2005 |
| 0.2132 | 2.215 | | unsexed | | ML | | China | Hainan Islands to Taiwan, 1997-1999 |
| 0.4946 | 2.217 | | unsexed | | ML | | China | South China Sea, 1997-1999 |

| | | | | | | | |
|------------------------|-------|-----|---------|------------|----|---------------|-------------------------|
| 0.2204 | 2.229 | | unsexed | | ML | China | Beibu Gulf, 1997-1999 |
| 0.1560 | 2.290 | Yes | female | 4.6 - 23.5 | ML | 32 Thailand | Andaman Sea / 2005-2005 |
| 0.1652 | 2.315 | No | female | 7.8 - 25.2 | ML | 223 Indonesia | Bangka Regency |

14) Length-length

(NA)

15) Length-frequencies

(NA)

16) Morphometrics

(NA)

17) Morphology

(NA)

18) Larvae

Larvae Information Summary for *Uroteuthis chinensis*

| | | | | |
|-----------------------------|-------------------------------|------------|------------|-------------|
| Main Ref: | Dong, Z. 1991 | | | |
| | Yolk-sac larvae | | | |
| | max | min | mod | Ref. |
| Length at birth (mm) | | | 0.3 | 75929 |
| Larval area | Unspecified (China) | | | |

19) Recruitment

(NA)

20) Abundance

(NA)

References

1. Roper, C.F.E., M.J. Sweeney and C.E. Nauen. 1984. (Ref. 275)
2. Arkhipkin, A. I., Rodhouse, P. G., Pierce, G. J., Sauer, W., Sakai, M., Allcock, L., Arguelles, J., Bower, J.R., Castillo, G., Ceriola, L., Chen, C-S., Chen, X., Diaz-Santana, M., Downey, N., Gonzales, A.F., Amores, J.G., Green, C.P., Guerra, A., Hendrickson, L.C., Ibanez, C. Ito, K., Jereb, P., Kato, Y.,

- Katugin, O.N., Kawano, M., Kidokoro, H., Kulik, V.V., Laptikhovsky, V.V., Lipinski, M.R., Liu, B., Mariategui, L., Marin, W., Medina, A., Miki, K., Miyahara, K., Moltschaniwakyj, N., Moustahfid, H., Nabhitabhata, J., Nanjo, N., Nigmatullin, C.M., Ohtani, T., Peci, G., Perez, J.A., Piatkowski, U., Saikliang, P., Salinas-Zavala, C.A., Steer, M., Tian, Y., Yamashiro, C., Tamashita, N. and Zeidberg, L.D., 2015. World Squid Fisheries. *Reviews in Fisheries Science and Aquaculture* 23(2): 92-252.
3. Chotiyaputta, C. 1995. Biology of cephalopods. In: Nabhitabhata, J. (ed.), *Biology and Culture of Cephalopods*, pp. 27-49. Rayong Coastal Aquaculture Station, Rayong.
 4. Chotiyaputta, C., Okutani, T. and Chaitiamyong, S. 1992. Systematic study of cephalopods in Thailand. Research Report National Research Council of Thailand, JSPS-NRCT International Cooperation Project. Bangkok, Thailand.
 5. IUCN. 2019. The IUCN Red List of Threatened Species. Version 2019-2. Available at: www.iucnredlist.org. (Accessed: 04 July 2019).
 6. Jereb, P., Piatkowski, U., Allcock, A.L., Belcari, P., Tasende, M.G., Gonzalez, A. Guerra, A., Hastie, L.C., Kafkadiou, E., Moreno, A., Pascual, S., Pereira, J., Pierce, G.J., Sanchez, P.S., Seixas, S., Sobrino, I. and Villanueva, R. 2010. Biology and ecology of cephalopod species commercially exploited in Europe: species accounts. In: Pierce, G.J., Allcock, L., Bruno, I., Bustamante, P., Gonzalez, A., Guerra, A., Jereb, P., Lefkaditou, E., Malham, S., Moreno, A., Pereira, J., Piatkowski, U., Raseo, M., Sanchez, P., Santos, M.B., Santurtun, M., Seixas, S., Sobrino, I., Villanueva, R. (ed.), *Cephalopod biology and fisheries in Europe. Co-operative Research Report 303.*, pp. 9-29. International Council for the Exploration of the Sea, Copenhagen.
 7. Jereb, P., Vecchione, M. and Roper, C.F.E. 2010. Family Loliginidae. In: P. Jereb and C.F.E. Roper (eds), *Cephalopods of the World. An annotated and illustrated catalogue of species known to date. Volume 2. Myopsid and Oegopsid squids*, pp. 38-117. FAO, Rome.
 8. Liao, C.H., Liu, T.-Y. and Hung, C.-Y. 2010. Morphometric variation between the swordtip (*Photololigo edulis*) and mitre (*P. chinensis*) squids in the waters off Taiwan. *Journal of Marine Science and Technology* 18(3): 405-412.
 9. Nateewathana, A. 1992. Taxonomic studies on loliginid squids (Cephalopoda: Loliginidae) from the Andaman Sea Coast of Thailand. *Phuket Marine Biological Center Research Bulletin* 57: 1-40.
 10. Songjitsawat, A. and Sookbuntoeng, S. 1988. Catch composition of marine fauna from light luring squid fishing. Technical paper no. 10. Eastern Marine Fisheries Development Center, Marine Fisheries Division, Department of Fisheries.
 11. Sweeney, M.J. 2012. Recent cephalopod primary type specimens: a searching tool. Available at: https://www.jiscmail.ac.uk/cgi-bin/filearea.cgi?LMGT1=FASTMOLL&a=get&f=/Type_Finder_Fastmoll-1.pdf.