


***Epinephelus coioides*****Orange-spotted grouper****Scientific classification**

Kingdom:	<a href="#">Animalia</a>
Phylum:	<a href="#">Chordata</a>
Class:	<a href="#">Actinopterygii</a>
Order:	<a href="#">Perciformes</a>
Family:	<a href="#">Serranidae</a>
Subfamily:	<a href="#">Epinephelinae</a>
Genus:	<a href="#">Epinephelus</a>
Species:	<b><i>E. coioides</i></b>

**Binomial name*****Epinephelus coioides*****([Hamilton](#), 1822)****A. Environment/Ecology:**

Marine; brackish; reef-associated; depth range 1 - 100 m (Ref. [167](#)). Subtropical; 37°N - 34°S, 28°E - 180°E (Ref. [5222](#))

**B. Distribution:**



Indo-West Pacific: Red Sea south to at least Durban, South Africa and eastward to Palau and Fiji, north to the Ryukyu Islands, south to the Arafura Sea (Ref. [9819](#)) and Australia. Recently reported from the Mediterranean coast of Israel (Ref. [5222](#)). Frequently misidentified as *Epinephelus tauvina* or *Epinephelus malabaricus* (Ref. [27362](#)).

#### C. Length at first maturity / Size / Weight / Age:

Maturity:  $L_m$  [48.3](#), range 25 - 30 cm Max length : 120 cm TL male/unsexed; (Ref. [47613](#)); max. published weight: 15.0 kg (Ref. [11228](#)); max. reported age: 22 years (Ref. [3627](#)) **Length at First Maturity= 25-30 cm Size= 55-75 Weight= 15kg Age= 2-3years (FiA,2020)**

#### D. Short description

**Dorsal spines** (total): 11; **Dorsal soft rays** (total): 13-16; **Anal spines**: 3; **Anal soft rays**: 8. This species is distinguished by the following characters: elongated body with greatest body depth at 2.9-3.7 in SL (for specimens 10-78 cm SL); head length 2.3-2.6 in SL. interorbital width 5.0-6.2 in HL; preopercle with enlarged serrae at angle and a broad shallow notch just above angle; upper edge of operculum straight or somewhat convex; maxilla reaches to or slightly past a vertical at rear edge of eye; upper jaw length 17-20% of SL; midlateral part of lower jaw with 2-3 rows of subequal teeth; gill rakers of first gill arch 8-10 + 14-17; pyloric caeca 50-60; lateral body scales rough, with minute auxiliary scales (body scales ctenoid except for nape, back, thorax, abdomen and above anal-fin base with cycloid scales); lateral-line scales 58-65; lateral-line tubes of anterior scales branched in adults. Colour: head and body tan dorsally, shading to whitish ventrally; numerous small brownish orange or reddish brown spots on head, body, and median fins; body with 5 faint, irregular, oblique, dark bars which bifurcate ventrally (irregular H-shaped bars); back with 3-4 blackish saddles; orange spots become poorly defined and darker with growth (Ref. [39231](#), [90102](#)).

#### E. Biology

Inhabit turbid coastal reefs (Ref. [9710](#)) and are often found in brackish water (Ref. [27362](#)) over mud and rubble (Ref. [6390](#)). Solitary (Ref 90102). Juveniles are common in shallow waters of estuaries over sand, mud and gravel and among mangroves (Ref. [6390](#)). Feed on small fishes, shrimps, and crabs. Probably spawn during restricted periods and form aggregations when doing so (Ref. [27352](#)). Females mature at 25 to 30 cm (2 to 3 years old), and sexual transition occurs at 55 to 75 cm (Ref. [39231](#)). Eggs and early larvae are probably pelagic (Ref. [6390](#)). Has been tested in several

countries as a potential species for mariculture (Ref. [43448](#)). Caught with hook-and-line, traps, trawls, and lift nets. Common and expensive in markets of the region; sold fresh and kept alive at restaurants in Asian countries (e.g. Hong Kong and Taiwan Province of China) (Ref. [39231](#)).

#### F. Life cycle and mating behavior

Pelagic spawner (Ref. [32184](#)).

#### G. Fisheries

(NA)

#### H. IUCN Red List Status

## GEOGRAPHIC RANGE

- **Taxonomy**

Kingdom:	<a href="#">Animalia</a>
Phylum:	<a href="#">Chordata</a>
Class:	<a href="#">Actinopterygii</a>
Order:	<a href="#">Perciformes</a>
Family:	<a href="#">Epinephelidae</a>
Genus:	<a href="#">Epinephelus</a>

- **Geographic Range**

**NUMBER OF LOCATIONS**

UPPER DEPTH LIMIT : 0 metres

LOWER DEPTH LIMIT : 100 metres

**RANGE DESCRIPTION**

This species is distributed in the Indo-Pacific Ocean from Durban, South Africa; north along East Africa, including Madagascar, Reunion and Mauritius, to the Red Sea and Persian Gulf; east to Palau and Fiji; north to the Ryukyu Islands, Japan; and south to the Arafura Sea and northern Australia (Heemstra and Randall 1993). It has also migrated through the Suez Canal to the eastern Mediterranean (Randall 1995). Its depth range is zero to 100 metres.

- **Population**

CURRENT POPULATION TREND : Decreasing

- **Habitat and Ecology**

System : Marine

Habitat type : Marine Neritic, Marine Intertidal


Generation Length : 12.5 years

- **Biological resource use :**

Fishing & harvesting aquatic resource

- **Threats**

Overfishing is a major threat to populations in the Persian Gulf (Grandcourt et al. 2005) and China, but this is not considered a major threat on a global-level at this time. Degradation of estuaries (juvenile habitat) and coral reefs (adult habitat) may also impact this species (Burke et al. 2002).



- **Use and Trade**

This species is important to fisheries through much of its range. It supplies live and chilled fish markets, both domestic and international. Mariculture of this species is extensive in Southeast Asia (Sadovy 2000). It is cultured for local consumption in Thailand and Singapore and for export using mainly wild-caught juvenile seed (Heemstra and Randall 1993, Yashiro 1996). Hatchery production has recently increased. Juveniles are still taken from the wild for grow-out in mariculture operations, but this is very poorly documented. This species is one of the most common next to *E. awoara* used to supply the large mariculture industry of Fujian, China; however, *E. coioides* juveniles are mainly sourced from hatchery production, not wild-caught fisheries (Liu and Sadovy 2009). Spawning aggregations are targeted by local fishers via spear in Papua New Guinea (Hamilton 2003). It is the most important reef-associated, commercial species in the Persian Gulf and is mainly collected via demersal traps (Grandcourt et al. 2005) as well as longlines and trawls (Carpenter et al. 1997). In northern Oman, fishermen target this species with semicircular wire basket traps (McIlwain et al. 2016). It is also commonly taken as bycatch in the shrimp and cutlassfish trawl fisheries of Iran (Raiesi et al. 2011, Paighambari and Daliri 2012, Hosseini et al. 2012).

- **Conservation Action**

Conservation measures directed to the Persian Gulf population have included reduction in fishing effort, modification of gear selectivity and the introduction of juvenile escape panels in the demersal trap fishery. In Qatar, fishery regulations include restricted mesh size, number of vessels, trip frequency and licensing (A. Al-Kuwayr pers. comm. 2013). However, these measures were found to be ineffective (Grandcourt et al. 2011). Suggested alternative measures include a moratorium on the use of traps in the offshore demersal fishery of Abu Dhabi (Grandcourt et al. 2011). There are no restrictions on fishing effort for this species in Oman (McIlwain et al. 2016). This species is under total protection in New South Wales, Australia (Public Consultation Document 2002) and there is a strict length limit enforced in Queensland, Australia (Fishing Industry Organization and Marketing

Amendment Regulation 1993). This species was released on Hong Kong artificial reefs in Yan Chau Tong and Hoi Ha Wan Marine Park in 2001 as a part of restocking trial (Cheung 2001), it is not known if this improved recovery.

I. More Information:

1) Stocks

(NA)

2) Ecology

<i>Ecology of Epinephelus coioides</i>	
<b>Main Ref.</b>	Randall, J.E. and P.C. Heemstra, 1991
<b>Distribution</b>	<p>Brackishwater</p> <ul style="list-style-type: none"> <li>• <b>estuaries/lagoons/brackish seas</b></li> <li>• <b>mangroves</b></li> </ul> <p>Highlighted items on the list are where <i>Epinephelus coioides</i> may be found.</p>
<b>Remarks</b>	Probably make frequent use of shelters, suggesting an 'ambush' method of feeding (Ref. 6390). Inhabit turbid coastal reefs (Ref. 9710) and are often found in brackish water (Ref. 27362) over mud and rubble (Ref. 6390). Juveniles are common in shallow waters of estuaries over sand, mud and gravel and among mangroves (Ref. 6390). Feed on small fishes, shrimps, and crabs. Probably spawn during restricted periods and form aggregations when doing so (Ref. 27352). Eggs and early larvae are probably pelagic (Ref. 6390).
<b>Substrate</b>	
<b>Substrate</b>	<b>Benthic; Soft Bottom:</b> mud; <b>Hard Bottom:</b> rubble;
<b>Substrate Ref.</b>	<a href="#">Letourneur, Y., P. Chabanet, P. Durville, M. Taquet, E. Teissier, M. Parmentier, J.-C. Quéro and K. Pothin, 2004</a>

3) Diet

Food and Feeding Habits: Diet Composition *Epinephelus coioides*

n = 1

Main Food	Percent	Trophic Level (y)	Predator Life Stage	Country	Locality	Ref.
<a href="#">nekton</a>	37	4.0	adults	New Caledonia	1985-1997	<a href="#">55797</a>
<a href="#">zoobenthos</a>	37	4.0	adults	New Caledonia	1985-1997	<a href="#">55797</a>

## 4) Reproduction

Reproduction of <i>Epinephelus coioides</i>	
Main Ref.	<a href="#">Heemstra, P.C. and J.E. Randall, 1993</a>
Mode	protogyny
Fertilization	external
Mating type	
Spawning frequency	one clear seasonal peak per year
Spawning aggregation	Yes. Ref. <a href="#">SCRFA, Science and Conservation of Fish Aggregations, 2018</a>
Batch spawner	Ref.
Reproductive guild	nonguarders open water/substratum egg scatterers
Parental Care	none
Description of life cycle and mating behavior	Pelagic spawner (Ref. 32184).
Search for more references on reproduction	<a href="#">Scirus</a>

## 5) Maturity

Maturity studies for <i>Epinephelus coioides</i>						
n = 5						
Sort by <input checked="" type="radio"/> Lm <input type="radio"/> Country <input type="radio"/> Locality <input type="radio"/> tm						
Lm (cm)	Length (cm)	Age range (y)	tm (y)	Sex of fish	Country	Locality
	49.0 -			<a href="#">unsexed</a>		Asia-Pacific Region
	25.0 - 30.0	2.0 - 3.0		<a href="#">female</a>		Arabian Gulf

43.5 TL	-	-	2.70	<a href="#">female</a>	United Arab Em.	coast of the Emirate of Abu Dhabi, Sept. 2000-March 2003
53.0 TL	-	-		<a href="#">female</a>	India	North Andhra Region (17°01'N-19°22'N; 83°23'E-85°14'E), Sept 2009-Oct 2011
92.0 TL	-	-		<a href="#">male</a>	India	North Andhra Region (17°01'N-19°22'N; 83°23'E-85°14'E), Sept 2009-Oct 2011

## 6) Spawning

*Spawning for **Epinephelus coioides***

n = 1

J	F	M	A	M	J	J	A	S	O	N	D	Country	Locality
		111	111	111								United Arab Em.	<a href="#">coast of the Emirate of Abu Dhabi, Sept. 2000-March 2003</a>

## 7) Spawning aggregation

*Spawning Aggregations of **Epinephelus coioides***

[ n = 1 ]

Country	Spawning type	Aggregation type	Status
<a href="#">Papua New Guinea</a>	Unknown	Unknown	Decreasing

## 8) Fecundity

*Fecundity for **Epinephelus coioides***

Sort by  Country  Locality

[ n = 1 ]

Country	Locality	Absolute Fecundity	
		min	max
India	<a href="#">North Andhra Region (17°01'N-19°22'N; 83°23'E-85°14'E), Sept 2009-Oct 2011</a>	43,618	463,940

## 9) Eggs

*Egg Characteristics of **Epinephelus coioides***

<b>Main Ref.</b>	<a href="#">Ordinio-Aguilar, R., H. Kohno, A. Ohno, M. Moteki and Y. Taki, 1995</a>
------------------	---

<b>Place of Development</b>	buoyant (pelagic)
<b>Shape of Egg</b>	spherical
<b>Attributes</b>	smooth, not sticky
<b>Color of Eggs</b>	transparent
<b>Get Information on</b>	<a href="#">Scirus</a>

10) Egg development

(NA)

11) Age/Size

List of Population Characteristics records for *Epinephelus coioides*  
n = 8

Sex	Wmax	Lmax (cm)	Tmax (y)	Country	Locality
<a href="#">unsexed</a>		95		Global	East Indies
<a href="#">unsexed</a>		97.9	12.2	United Arab Em.	Abu Dhabi, 2002-2003
<a href="#">unsexed</a>	1.5 kg	100		South Africa	Not specified
<a href="#">unsexed</a>		100		Indonesia	Not specified
<a href="#">unsexed</a>		111	22	Kuwait	Kuwait
<a href="#">unsexed</a>		120			Asia-Pacific Region
<a href="#">unsexed</a>		120		Oman	Not specified
<a href="#">unsexed</a>		120		Iran	Persian Gulf and Oman Sea

12) Growth

**Growth parameters for *Epinephelus coioides***

Maximum Length 120cm TL

n = 2

Note that studies where L<sub>oo</sub> is very different (+/- 1/3) from L<sub>max</sub> are doubtful.

<a href="#">M vs K graph</a>	[n = 1]
<a href="#">M vs L<sub>inf</sub> graph</a>	[n = 1]
<a href="#">Longevity vs 3/K graph</a>	[n = 1]

$\phi = 3.13$  L<sub>inf</sub> = 97.9 cm TL K = 0.1



Median record no. 2

55546Ref. [55546](#)

Length (cm)	Length Type	K (1/y)	to (years)	M (1/y)	Temp° C	Ø'	Country	Locality	Questionable	Captive
93.0	TL	<a href="#">0.167</a>		0.20	23.5	3.16	Kuwait		No	No
97.9	TL	<a href="#">0.140</a>	-1.50			3.13	United Arab Em.	Abu Dhabi	No	No

## 13) Length-weight

Length-Weight Parameters for <i>Epinephelus coioides</i>									
<a href="#">Length-weight (log a vs b) graph</a>							[n=6]		
							<a href="#">Hide graph</a>		
Sort by <input type="radio"/> a <input checked="" type="radio"/> b <input type="radio"/> Country <input type="radio"/> Locality									
Score	a	b	Sex	Length (cm)	Length type	r <sup>2</sup>	n	Country	Locality
0.96	<a href="#">0.01990</a>	2.990	mixed	14.2 - 107.0	TL	0.956	440	India	Visakhapatnam, Andhra Pradesh / 2009-2011
0.60	<a href="#">0.01440</a>	3.024	unsexed		TL		1912	Kuwait	
1.00	<a href="#">0.02160</a>	3.053	unsexed	16.0 - 51.0	SL	0.995	15	Philippines	Davao Gulf / 2009-2012
0.99	<a href="#">0.01050</a>	3.084	unsexed	6.5 - 111.0	TL	0.994	41	New Caledonia	lagoon
0.99	<a href="#">0.00990</a>	3.102	mixed	6.5 - 111.0	FL	0.994	44	New Caledonia	
0.50	<a href="#">0.01020</a>	3.232	Mixed		TL			United Arab Em.	Abu Dhabi, 2002-2003

## 14) Length-length

Length-length Parameters for <i>Epinephelus coioides</i>						
[n=6]						
Unknown length	a	b	Known length	r	Length range (cm)	Sex of fish
<a href="#">SL</a>	-0.630	0.840	TL	0.999773	11.2 - 82.5	unsexed
<a href="#">TL</a>	0.000	1.000	FL		-	unsexed
<a href="#">TL</a>	0.000	1.148	SL		-	unsexed
<a href="#">TL</a>	0.000	1.217	SL		-	unsexed
<a href="#">TL</a>	0.349	1.223	SL	0.998	16 - 51	unsexed

<a href="#">TL</a>	0.000	1.227	SL		-	unsexed
--------------------	-------	-------	----	--	---	---------

## 15) Length-frequencies

(NA)

## 16) Morphometrics

Morphometric Data for *Epinephelus coioides*

n = 4

Picture Name	Length		Lifestage	Aspect ratio
<a href="#">Epcoi_u0.gif</a>	31.9	SL	unsexed	0.99
<a href="#">Epcoi_u1.jpg</a>	9.1	SL	juvenile	1.42
<a href="#">Epcoi_u3.jpg</a>	12.9	SL	unsexed	1.25
<a href="#">Epcoi_u5.jpg</a>	19.6	SL	unsexed	

Picture Used	EPCOI_U0.GIF
Size (cm)	31.9 SL
Sex	unsexed
Total length (TL)	584 pixels
Standard length	82.2 % TL
Fork length	100.0 % TL
Pre-anal length	59.9 % TL
Pre-dorsal length	32.4 % TL
Pre-pelvic length	30.5 % TL
Pre-pectoral length	32.0 % TL
Body depth	23.8 % TL
Head length (HL)	35.4 % TL
Eye diameter	11.6 % HL
Pre-orbital length	23.7 % HL
Aspect ratio of caudal fin	0.992808

Picture Used	Epcoi_u1.jpg
Size (cm)	9.1 SL, 11.2
Sex	juvenile
Total length (TL)	605 pixels
Standard length	87.1 % TL
Fork length	100.0 % TL

Pre-anal length	57.2 % TL
Pre-dorsal length	28.1 % TL
Pre-pelvic length	30.6 % TL
Pre-pectoral length	29.1 % TL
Body depth	26.3 % TL
Head length (HL)	32.4 % TL
Eye diameter	19.9 % HL
Pre-orbital length	18.4 % HL
Aspect ratio of caudal fin	1.41858

Picture Used	EPCOI_U3.JPG
Size (cm)	12.9 SL, 16.3
Sex	unsexed
Total length (TL)	584 pixels
Standard length	81.5 % TL
Fork length	100.0 % TL
Pre-anal length	56.2 % TL
Pre-dorsal length	28.9 % TL
Pre-pelvic length	29.6 % TL
Pre-pectoral length	26.9 % TL
Body depth	25.7 % TL
Head length (HL)	31.8 % TL
Eye diameter	18.3 % HL
Pre-orbital length	24.2 % HL
Aspect ratio of caudal fin	1.25261

Picture Used	Epcoi_u5.jpg
Size (cm)	19.6 SL, 24.9
Sex	unsexed
Total length (TL)	600 pixels
Standard length	83.3 % TL
Fork length	100.0 % TL
Pre-anal length	55.0 % TL
Pre-dorsal length	27.3 % TL
Pre-pelvic length	28.5 % TL

Pre-pectoral length	27.5 % TL
Body depth	24.5 % TL
Head length (HL)	29.8 % TL
Eye diameter	13.4 % HL
Pre-orbital length	22.9 % HL
Aspect ratio of caudal fin	1.47271

## 17) Morphology

(NA)

## 18) Larvae

Larvae Information Summary for *Epinephelus coioides***Main Ref:**

[Kailola, P.J., M.J. Williams, P.C. Stewart, R.E. Reichelt, A. McNee and C. Grieve 1993](#)

Yolk-sac larvae

**Place of development**

planktonic

**Larval area**

North West Shelf, Australia

## 19) Recruitment

(NA)

## 20) Abundance

(NA)

## References

1. Heemstra, P.C. and J.E. Randall, 1993. FAO Species Catalogue. Vol. 16. Groupers of the world (family Serranidae, subfamily Epinephelinae). An annotated and illustrated catalogue of the grouper, rockcod, hind, coral grouper and lyretail species known to date. Rome: FAO. FAO Fish. Synop. 125(16):382 p. (Ref. [5222](#))
2. Agembe, S., Mlewa, C.M. and Arara-Kaunda, B. 2010. Catch composition, abundance and length-weight relationships of groupers (Pisces: Serranidae) from inshore waters of Kenya. *Western Indian Ocean Journal of Marine Science* 9: 91-102.

3. Anonymous. 1993. Fishing Industry Organization and Marketing Amendment Regulation (No. 3). Subordinate Legislation No. 235. State of Queensland.
4. Anonymous. 2002. Public Consultation Document. Environmental Impact Statement on the Ocean Hauling Fishery - Appendix F. Department of the Environment and Heritage, Canberra, Australia
5. Burke, L., Selig, E. and Spalding, M. 2002. *Reefs at Risk in Southeast Asia*. World Resources Institute.
6. Carpenter, K.E., Harrison, P.L., Hodgson, G., Alsaffer, A.H. and Alhazeem, S.H. 1997a. *The Corals and Coral Reef Fishes of Kuwait*. Kuwait Institute for Scientific Research and the Environmental Public Authority, Safat, Kuwait.
7. Cheung, W.L. 2001. Restocking: an effective measures to restore the depleted fishery stocks in Hong Kong. *Porcupine* 24: 12-14.
8. Depczynski, M. and Bellwood, D.R. 2006. Extremes, plasticity, and invariance in vertebrate life history traits: insights from coral reef fishes. *Ecology* 87: 3119-3127.
9. Feary, D.A., Burt, J.A. and Bartholomew, A. 2011. Artificial marine habitats in the Arabian Gulf: Review of current use, benefits and management implications. *Ocean and Coastal Management* 54: 742-749.
10. Grandcourt, E.M., Al Abdessalaam, T.Z., Francis, F., Al Shamsi, A.T. and Hartmann, S.A. 2009. Reproductive biology and implications for management of the orange-spotted grouper *Epinephelus coioides* in the southern Arabian Gulf. *Journal of Fish Biology* 74: 820-841.
11. Grandcourt E.M., Al Abdessalaam T.Z., Francis F. and Al Shamsi, A.T. 2003. Population biology and stock assessment of the Orange-spotted grouper, *Epinephelus coioides*, in the Southern Arabian Gulf. Internal report, Marine Environmental Research Centre, Environmental Research and Wildlife Development Agency, UAE.
12. Grandcourt, E.M., Al Abdessalaam, T.Z., Francis, F. and Al Shamsi, A.T. 2005. Population biology and assessment of the orange-spotted grouper, *Epinephelus coioides* (Hamilton, 1822), in the southern Arabian Gulf. *Fisheries Research* 74: 55-68.
13. Grandcourt, E.M., Al Abdessalaam, T.Z., Hartmann, S.A., Al Shamsi, A.T. and Francis, F. 2011b. An Evaluation of the Selectivity Characteristics of Different Juvenile Fish Escape Panel Designs for the Demersal Trap Fishery of Abu Dhabi, United Arab Emirates. *Open Journal of Marine Science* 3: 98-107.
14. Grandcourt, E.M., Al Abdessalaam, T.Z., Hartmann, S.A., Francis, F. and Al Shamsi, A.T. 2011a. An Evaluation of the Management Effectiveness of a Stock Re-Building Strategy for a Multi-Species Demersal Trap Fishery in Abu Dhabi, United Arab Emirates. *Open Journal of Marine Science* 3: 82-97.
15. Hamilton, R. 2003. A report on the status of exploited reef fish aggregations in the Solomon Islands and Papua New Guinea- Choiseul, Ysabel, Bougainville and Manus Provinces. Western Pacific Fisher Survey Series. Society for the Conservation of Reef Fish Aggregations.
16. Heemstra, P.C. and Randall, J.E. 1993. *FAO species catalogue. Vol. 16. Groupers of the world (Family Serranidae, Subfamily Epinephelinae). An annotated and illustrated catalogue of the grouper, rockcod, hind, coral grouper and lyretail species known to date*. FAO Fisheries Synopsis. No. 125, Vol. 16. Rome, FAO.
17. Hosseini, S.A., Raeisi, H. and Paighambari, S.Y. 2012. Temporal and Spatial Variations of Finfish Bycatch of Cutlassfish Trawl in Bushehr and Hormozgan Marine Waters, the Northern Persian Gulf. *Journal of the Persian Gulf* 3(9): 1-8.
18. IUCN. 2018. The IUCN Red List of Threatened Species. Version 2018-2. Available at: [www.iucnredlist.org](http://www.iucnredlist.org). (Accessed: 15 November 2018).
19. Kailola, P.J., Williams, M.J., Stewart, P.C., Reichelt, R.E., McNee, A. and Grieve, C. 1993. *Australian fisheries resources*. Bureau of Resource Sciences, Canberra, Australia.

20. Kandula, S., Shrikanya, K. V. and Iswarya Deepti, V. A. 2015. Species diversity and some aspects of reproductive biology and life history of groupers (Pisces: Serranidae: Epinephelinae) off the central eastern coast of India. *Marine Biology Research* 11(1): 18-33.
21. Lam, S.S.T. 2013. Status of Live Reef Fish Trade in the Hong Kong Retail Sector. Unpublished Final Year Project, The University of Hong Kong.
22. Liu, M. and Mitcheson, Y. 2011. The influence of social factors on juvenile sexual differentiation in a diandric, protogynous grouper *Epinephelus coioides*. *Ichthyological Research* 58: 84-89.
23. Mathews, C.P. and Samuel, M. 1991. Growth, mortality and length-weight parameters for some Kuwaiti fish and shrimp. *Fishbyte* 9((2):): 30-33.
24. McIlwain, J., Ambu-ali, A., Jardani, N., Halford, A., Al-Oufi, H. and Feary, D. 2015. Demographic profile of an overexploited serranid, the orange-spotted grouper (*Epinephelus coioides*), from northern Oman. *Fish Bulletin NOAA* 114: 490-502.
25. Mous, P.J. and Pet, J. 2016. Length-based assessment of data-poor multi-species dpppe grouper fisheries in fisheries management areas (WPP) 573,712, 713, 714, 715 & 718 in Indonesia. TNC-IPCP Technical Paper (unpublished draft). The Nature Conservancy, Indonesia.
26. Paighambari, S.Y. and Daliri, M. 2012. The bycatch composition of shrimp trawl fisheries in Bushehr coastal waters, the northern Persian Gulf. *Journal of the Persian Gulf* 3(7): 27-36.
27. Pauly, D. and Zeller, D. 2015. Sea Around Us Concepts, Design and Data. Available at: [seaaroundus.org](http://seaaroundus.org).
28. Raeisi, H., Hosseini, S.A., Paighambari, S.Y., Taghavi, S.A.A. and Davoodi, R. 2011. Species composition and depth variation of cutlassfish (*Trichiurus lepturus* L. 1785) trawl bycatch in the fishing grounds of Bushehr waters, Persian Gulf. *African Journal of Biotechnology* 10(76): 17610-17619.
29. Randall, J.E. 1995. *Coastal fishes of Oman*. University of Hawaii Press, Honolulu, Hawaii.
30. Randall, J.E., Allen, G.R. and Steene, R.C. 1997. *Fishes of the Great Barrier Reef and Coral Sea*. Revised edition. Crawford House, Bathurst, NSW. 557pp.
31. Sadovy, Y. 2000. Regional survey for fry/fingerling supply and current practices for grouper mariculture: evaluating current status and long-term prospects for grouper mariculture in South East Asia. Final report to the Collaborative APEC grouper research and development network (FWG 01/99). December, 2000.
32. Sadovy, Y. and Cornish, A.S. 2000. *Reef Fishes of Hong Kong*. Hong Kong University Press, Hong Kong. 320 pp.
33. Shallard, B. and Associates. 2003. Fish Resource Assessment Survey Project of Abu Dhabi and UAE Waters: Project Report. Environmental Research and Wildlife Development Agency Abu Dhabi, Abu Dhabi.
34. Shao, K. H. S. and Ho, K. Y. 2016. Rapid Survey on the Species Composition and Spatial Patterns of Reef Fishes Associated with Shallow Coral Communities, Rocky Reefs and Adjacent Areas in Hong Kong Waters 2016 – 2017. Report submitted to The Swire Group Charitable Trust, Unpublished Report.
35. Shea, K.H.S. and Ho, K.Y. 2016. Rapid survey on the species composition and spatial patterns of reef fishes associated with shallow coral communities, rocky reefs and adjacent areas in Hong Kong waters 2016 – 2017. Unpublished report submitted to the Swire Group Charitable Trust.
36. Sluka, R.D. and Lazarus, S. 2010. Grouper (Pisces: Serranidae) relative abundance and diversity on the west coast of India. *Marine Biodiversity Records* 3: 1-3.
37. To, A.W.L. and Sadovy de Mitcheson, Y. 2009. Shrinking baseline: the growth in juvenile fisheries, with the Hong Kong grouper fishery as a case study. *Fish and Fisheries* 10(4): 396-407.
38. To, W. 2009. The biology, fishery of groupers (family: Serranidae) in Hong Kong and adjacent waters, and implications for management. University of Hong Kong.

39. Valinassab, T., Daryanabard, R., Dehghani, R. and Pierce, G.J. 2006. Abundance of demersal fish resources in the Persian Gulf and Oman Sea. *Journal of the Marine Biological Association of the United Kingdom* 86: 1455-1462.
40. Yashiro, R. 1996. Status of Grouper Breeding and Culture in Thailand. Online report (Accessed: 30/04/2004).