

Rastrelliger brachysoma**Indo-Pacific Mackerel or Short Mackerel****Scientific classification**

Kingdom: [Animalia](#)
Phylum: [Chordata](#)
Class: [Actinopterygii](#)
Order: [Scombriformes](#)
Suborder: [Scombroidei](#)
Family: [Scombridae](#)
Subfamily: [Scombrinae](#)
Tribe: [Scombrini](#)
Genus: [Rastrelliger](#)
Species: [R. brachysoma](#)

Binomial name

Rastrelliger brachysoma

([Bleeker](#), 1851)

Synonyms


- *Rastrelliger neglectus* van Kampen,
1907
- *Scomber brachysoma* Bleeker, 1851

- *Scomber neglectus* van Kampen,
1907

A. Environment/Ecology:

Marine; brackish; pelagic-neritic; oceanodromous (Ref. [51243](#)); depth range 15 - 200 m (Ref. [28016](#)). Tropical; 20°C - 30°C (Ref. [54858](#)); 18°N - 18°S, 93°E - 180°E (Ref. [54858](#))

B. Distribution:



Pacific Ocean: Andaman Sea to Thailand, Indonesia, Papua New Guinea, Philippines, Solomon Islands and Fiji.

C. Length at first maturity / Size / Weight / Age:

Maturity: L_m [17.0](#) range ? - ? cm **Max length :** 34.5 cm FL male/unsexed; (Ref. [168](#)); common length : 20.0 cm FL male/unsexed; (Ref. [168](#)) **Length at First Maturity=** total length of 16.83 for male and 17.18 cm for female **Size=** maximize size of 17.15cm for male and 17.70 cm for female **Weight=** 55.05g for male and 58.01 for female **Age=** ??(FiA,2020). **Length at First Maturity=**16.98 cm L_{inf} = 22.23, common length = 16-18 cm(Srichanngam et al.,2014)

D. Short description

Dorsal spines (total): 8 - 11; **Dorsal soft rays** (total): 12; **Anal spines**: 0; **Anal soft rays**: 12; **Vertebrae**: 31. This species is distinguished by the following characters: body very deep, its depth at posterior margin of opercle 3.7-4.3 times in fork length; head equal to or less than body depth; maxilla covered by lacrimal bone but extending nearly to end of lacrimal; gill rakers very long, visible when mouth is opened, 30-48 on lower limb of first gill arch; numerous bristles on longest gill raker, about 150 on one side in specimens of 12.7 cm, 210 in specimens of 16 cm, and 240 at 19 cm fork length; intestine very long, 3.2-3.6 times fork length; snout pointed; swim bladder present; vertebrae 13 +

18 = 31; interpelvic process small and single; anal spine rudimentary. Colour of spinous dorsal fin yellowish with a black edge, pectoral and pelvic fins dusky, other fins yellowish (Ref. [168](#), [9684](#)).

E. Biology

An epipelagic, neritic species that tolerates slightly reduced salinities in estuarine habitats and in areas where surface temperature range between 20° and 30°C. It forms schools of equally sized individuals. Batch spawning is believed to extend from March through September. Feeds chiefly on microzooplankton with a high phytoplankton component. Marketed fresh, frozen, canned, dried salted and smoked (Ref. [168](#), [9684](#)).

F. Life cycle and mating behavior

G. Fisheries

Catches of this species are usually either recorded as *Rastrelliger* spp. or are combined with *R. kanagurta*. It is the most important commercial species of mackerel in the Philippines, caught the year round with native purse seines (italakop) and fish corrals (ibaklad) in Manila Bay (Manacop, 1958) and by dynamiting. In India, ("indian mackerel fishing"), it is fished with a variety of gear such as gillnets, seines , and cast nets and drift nets operated from boats with out-riggers and from dugout canoes. The catch in the Philippines fluctuated between 25 183 metric tons in 1978 and 18 962 metric tons in 1981 (FAO, 1983). The total catch reported for this species to FAO for 1999 was 25 713 t. The countries with the largest catches were Philippines (25 713 t).

H. IUCN Red List Status

GEOGRAPHIC RANGE

- **Taxonomy**

| | |
|----------|--------------------------------|
| Kingdom: | Animalia |
| Phylum: | Chordata |
| Class: | Actinopterygii |
| Order: | Perciformes |
| Family: | Scombridae |
| Genus: | Rastrelliger |

- **Geographic Range**

NUMBER OF LOCATIONS

UPPER DEPTH LIMIT : 0 metres

LOWER DEPTH LIMIT : 200 metres

RANGE DESCRIPTION

This species is found in the Pacific Ocean from the Andaman Sea to Thailand, Indonesia, Papua New Guinea, Philippines, Solomon Islands and Fiji.

- Population**

CURRENT POPULATION TREND : Unknown
POPULATION SEVERELY FRAGMENTED : No

- Habitat and Ecology**

System : Marine
Habitat type : Marine Neritic
Movement patterns : Full Migrant


- Biological resource use :**

Logging & wood harvesting

- Threats**

This species is highly commercial, and is caught with a number of different gears including purse-seines, fish corrals, gill-nets, cast and drift nets, and by dynamiting. It is marketed fresh, frozen, canned, dried salted and smoked (Collette 2001). In the Philippines, this is a highly commercial species caught by seines, and where landings range from 10,000 to 50,000 tonnes per year.

Worldwide reported landings of *Rastrelliger* spp. are increasing, and although there is no information on effort, it is also assumed to be increasing.



- Use and Trade**

This is a highly commercial fish species.

- Conservation Action**

There are no species-specific conservation measures. Although landings are increasing, without information on effort, it is not known if current fishing activities are affecting population abundance. Better reporting is needed to determine species specific landings if possible. Additionally, given the high combined landings for this species and unknown level of effort and the absence of an international management body, further monitoring of this species is needed on the national level.

I. More Information:

1) Stocks

This species is widespread in southeastern Asia. There is no information on population or general abundance. This species is targeted in commercial and artisanal fisheries throughout its range, but landings are primarily reported in combination with mixed *Rastrelliger* spp. Reported worldwide landings for *Rastrelliger* species have steadily increased since 1950 to over 800,000 tonnes in 2006, but no effort information is available. Given that effort is assumed to be increasing, it is not known how this species population is affected by current and historical fishing pressure. This species is listed as Data Deficient. Given the absence of an international management body, further monitoring of this species is needed on the national level, in addition to species-specific data on landings, effort and population status.

Catches of the three species of *Rastrelliger* are not usually recorded separately. *Rastrelliger brachysoma* is the most important commercial species of mackerel in the Philippines (Collette and Nauen 1983). Reported worldwide landings show gradual increase for all three *Rastrelliger* species, with combined reported landings increasing from 200,000 tonnes in 1950 to over 800,000 tonnes in 2006 (FAO 2009).

2) Ecology

Ecology of *Rastrelliger brachysoma*

This species is pelagic and oceanodromous and is found in estuarine habitats with slightly reduced salinities and in areas where surface temperature range between 20–30°C. It forms schools of equally sized individuals, and feeds chiefly on microzooplankton with a high phytoplankton component.

| | | |
|--|--|---|
| Main Ref. | Collette, B.B. and C.E. Nauen, 1983 | |
| Distribution | Marine - Oceanic <ul style="list-style-type: none"> • epipelagic | Brackishwater <ul style="list-style-type: none"> • estuaries/lagoons/brackish seas |
| Highlighted items on the list are where <i>Rastrelliger brachysoma</i> may be found. | | |
| Remarks | This species tolerates slightly reduced salinities in estuarine habitats. Feeds chiefly on microzooplankton with a high phytoplankton component. | |

3) Diet

Feeding

The short mackerel feeds chiefly on microzooplankton with a high phytoplankton component.

| | |
|------------------|---|
| Feeding type | plants/detritus+animals (troph. 2.2-2.79) |
| Feeding type ref | Collette, B.B. and C.E. Nauen, 1983 |

| | | | | | | | |
|--|-----------------------------------|---|------|------------------------|----------------------------|--|--|
| | Feeding habit | selective plankton feeding | | | | | |
| | Feeding habit ref | Collette, B.B. and C.E. Nauen, 1983 | | | | | |
| | Trophic level(s) | | | Original sample | Unfished population | Remark | |
| | Estimation method | Troph | s.e. | Troph | s.e. | | |
| | From individual food items | | | | | Trophic level estimated from a number of food items using a randomized resampling routine. | |
| | | 2.72 | 0.31 | | | | |

4) Reproduction

| Reproduction of <i>Rastrelliger brachysoma</i> | |
|--|---|
| Main Ref. | Collette, B.B. and C.E. Nauen, 1983 |
| Mode | dioecism |
| Fertilization | external |
| Mating type | |
| Spawning frequency | |
| Spawning aggregation | Ref. |
| Batch spawner | Yes. Ref. Collette, B.B. and C.E. Nauen, 1983 |
| Reproductive guild | nonguarders open water/substratum egg scatterers |
| Parental Care | none |
| Description of life cycle and mating behavior | |
| Search for more references on reproduction | Scirus |

5) Maturity

| Maturity studies for <i>Rastrelliger brachysoma</i> n = 1. (Lm vs Linf graph) | | | | | | | |
|---|-------------|---------------|--------|-------------------------|----------|------------------|--|
| Lm (cm) | Length (cm) | Age range (y) | tm (y) | Sex of fish | Country | Locality | |
| 17.0 | - | - | - | unsexed | Thailand | Gulf of Thailand | |

6) Spawning

| Spawning for <i>Rastrelliger brachysoma</i> | | | | | | | | | | | | | |
|--|---|---|-----|-----|-----|-----|-----|-----|-----|-----|---|---------|----------------------------------|
| n = 1 | | | | | | | | | | | | | |
| J | F | M | A | M | J | J | A | S | O | N | D | Country | Locality |
| | | | 111 | 111 | 111 | 111 | 111 | 111 | 111 | 111 | | | <u>Central Indo-West Pacific</u> |

7) Spawning aggregation

Batch spawning is believed to extend from March through September.

8) Fecundity

(NA)

9) Eggs

(NA)

10) Egg development

(NA)

11) Age/Size

| List of Population Characteristics records for <i>Rastrelliger brachysoma</i> | | | | | |
|--|------|-----------|----------|-------------|-----------------------------------|
| n = 7 | | | | | |
| Sex | Wmax | Lmax (cm) | Tmax (y) | Country | Locality |
| <u>unsexed</u> | | | 2 | Thailand | Gulf of Thailand |
| <u>unsexed</u> | | 13.2 | | Philippines | Estancia, Iloilo |
| <u>unsexed</u> | | 22 | | Malaysia | Kedah |
| <u>unsexed</u> | | 22 | | Indonesia | Sumatra |
| <u>unsexed</u> | | 24 | | Thailand | Gulf of Thailand, 10° N 100° E |
| <u>unsexed</u> | | 34 | | | to be filled |
| <u>unsexed</u> | | 35 | | Philippines | Not specified |

12) Growth

Growth parameters for *Rastrelliger brachysoma*

Maximum Length 34.5cm FL n = 38

Note that studies where L_{oo} is very different (+/- 1/3) from L_{max} are doubtful.

L_m vs L_{inf} graph [n = 2]

Reproductive graph [n = 1]

M vs K graph [n = 2]

M vs L_{inf} graph [n = 2]

Longevity vs 3/K graph [n = 1]

$\phi = 3.00$ $L_{inf} = 25.0$ cm FL $K = 1.6$ Median record no. 201263 Ref. [1263](#)

| L _{oo} (c m) | Lengt h Type | K (1/ y) | to (year s) | Se x | M (1/ y) | Temp° C | L m | ϕ ' | Countr y | Locality | Questiona ble | Capti ve |
|-----------------------------|--------------------|-----------------------|-------------------|---------|----------------|------------|----------|-------------|-------------|--------------------------------|------------------|-------------|
| 18.2 | TL | 1.560 | | | | 28.0 | | 2.7 1 | Thailand | Gulf of Thailand | No | No |
| 19.6 | TL | 4.140 | | | | 28.0 | | 3.2 0 | Thailand | 10°N 100°E Gulf of Thailand | No | No |
| 20.0 | TL | 3.528 | | | | 28.0 | | 3.1 5 | Thailand | 10°N 100°E Gulf of Thailand | No | No |
| 20.9 | TL | 3.384 | 0.00 | | 7.22 | 28.0 | | 3.1 7 | Thailand | Gulf of Thailand | No | No |
| 20.9 | TL | 4.200 | | | | 28.0 | | 3.2 6 | Thailand | Inner Gulf of Thailand | No | No |
| 21.2 | FL | 0.960 | | | | 23.0 | | 2.6 3 | Philippines | Samar Sea | No | No |
| 22.0 | TL | 0.700 | | | | 28.0 | 17. 0 | 2.5 3 | Thailand | Gulf of Thailand | No | No |
| 22.0 | SL | 1.420 | | | | 21.5 | | 2.8 4 | Thailand | Southwest coast | No | No |
| 22.4 | TL | 2.000 | | | | 21.5 | | 3.0 0 | Thailand | Strait of Malacca | No | No |
| 22.9 | TL | 2.280 | | | 4.56 | 28.0 | | 3.0 8 | Indonesia | Tanjung Satai (Western Borneo) | No | No |
| 22.9 | TL | 1.800 | | | | 27.0 | | 2.9 7 | Indonesia | Java Sea (Central Java) | No | No |

| | | | | | | | | | | | |
|------|----|--------------|--|--|------|----------|----------|-------------|-----------------------|----|----|
| 23.0 | TL | <u>3.600</u> | | | 28.0 | | 3.2 8 | Thailand | Gulf of Thailand | No | No |
| 23.2 | FL | <u>1.200</u> | | | 23.0 | | 2.8 1 | Philippines | Guimaras Strait | No | No |
| 23.5 | TL | <u>1.500</u> | | | 29.0 | | 2.9 2 | Malaysia | Kedah | No | No |
| 24.0 | TL | <u>1.020</u> | | | 22.4 | | 2.7 7 | Malaysia | Selangor | No | No |
| 24.0 | TL | <u>1.040</u> | | | 29.0 | | 2.7 8 | Malaysia | Kedah | No | No |
| 24.2 | TL | <u>0.520</u> | | | 22.4 | | 2.4 8 | Malaysia | Selangor | No | No |
| 24.5 | FL | <u>1.280</u> | | | 28.5 | | 2.8 9 | Philippines | Ragay Gulf | No | No |
| 24.5 | TL | <u>1.400</u> | | | 21.5 | | 2.9 2 | Thailand | Strait of Malacca | No | No |
| 25.0 | FL | <u>1.600</u> | | | 28.5 | | 3.0 0 | Philippines | Samar Sea | No | No |
| 25.0 | FL | <u>1.300</u> | | | 28.5 | | 2.9 1 | Philippines | Samar Sea | No | No |
| 25.0 | TL | <u>0.820</u> | | | 22.4 | | 2.7 1 | Malaysia | Perak | No | No |
| 25.1 | TL | <u>1.250</u> | | | 21.5 | | 2.9 0 | Thailand | Strait of Malacca | No | No |
| 25.4 | TL | <u>1.330</u> | | | 21.5 | | 2.9 3 | Thailand | Strait of Malacca | No | No |
| 25.5 | FL | <u>1.450</u> | | | 28.5 | | 2.9 7 | Philippines | Samar Sea | No | No |
| 25.8 | TL | <u>1.630</u> | | | 28.0 | | 3.0 4 | Indonesia | Java Sea (Pekalongan) | No | No |
| 26.0 | TL | <u>0.600</u> | | | 22.4 | | 2.6 1 | Malaysia | Perak | No | No |
| 26.3 | TL | <u>1.300</u> | | | 21.5 | | 2.9 5 | Thailand | Strait of Malacca | No | No |
| 26.5 | TL | <u>1.050</u> | | | 12.0 | | 2.8 7 | Indonesia | Asahan, Sumatra | No | No |
| 27.0 | FL | <u>1.600</u> | | | 27.0 | | 3.0 7 | Myanmar | Mergui Archipelago | No | No |
| 28.0 | | <u>1.000</u> | | | | 11. 9 | 2.8 9 | Philippines | Davao Gulf | No | No |

| | | | | | | | | | | | |
|------|----|--------------|--|--|------|--|----------|-------------|--------------------------|----|----|
| 28.5 | FL | <u>1.400</u> | | | 28.0 | | 3.0 6 | Philippines | Guimaras Strait | No | No |
| 29.8 | TL | <u>1.300</u> | | | 28.5 | | 3.0 6 | Philippines | Samar Sea | No | No |
| 32.5 | TL | <u>1.200</u> | | | 27.7 | | 3.1 0 | Philippines | Visayan Sea | No | No |
| 34.0 | TL | <u>1.100</u> | | | 28.2 | | 3.1 0 | Philippines | Manila Bay | No | No |
| 34.0 | TL | <u>0.980</u> | | | 28.2 | | 3.0 5 | Philippines | Manila Bay | No | No |
| 34.0 | TL | <u>0.981</u> | | | 27.7 | | 3.0 5 | Philippines | Visayan Sea, 1983 & 1985 | No | No |
| 34.0 | TL | <u>0.982</u> | | | 28.4 | | 3.0 6 | Philippines | Leyte Gulf | No | No |

13) Length-weight

Length-Weight Parameters for *Rastrelliger brachysoma*

| Score | a | b | Sex | Length (cm) | Length type | r ² | Country | Locality |
|-------|----------------|-------|---------|-------------|-------------|----------------|-------------|-------------------------------|
| 1.00 | <u>0.02580</u> | 2.879 | unsexed | | | 1.000 | Indonesia | Tanjung Satai, Western Borneo |
| 0.50 | <u>0.01300</u> | 3.210 | unsexed | | | | Thailand | Indian coast, 1967-77 |
| 0.50 | <u>0.00614</u> | 3.213 | unsexed | | | | Thailand | Inner Gulf of Thailand |
| 0.50 | <u>0.01000</u> | 3.230 | unsexed | | | | Philippines | Guimaras Strait, 1988-89 |

14) Length-length

Length-Length Parameters for *Rastrelliger brachysoma*

n=4

| Unknown length | a | b | Known length | r | Length range (cm) | Sex of fish |
|----------------|-------|-------|--------------|---|-------------------|-------------|
| <u>SL</u> | 0.000 | 0.901 | FL | | - | unsexed |
| <u>SL</u> | 0.000 | 0.961 | FL | | - | unsexed |

| | | | | | | |
|-----------|-------|-------|----|--|---|---------|
| <u>TL</u> | 0.000 | 1.104 | FL | | - | unsexed |
| <u>TL</u> | 0.000 | 1.149 | SL | | - | unsexed |

15) Length-frequencies

List of frequency studies for *Rastrelliger brachysoma*

| Locality | Year from - to | Sex | Gear | Frequency type |
|--|----------------|---------------|---------------|--------------------------|
| Guimaras Strait, Philippines | 1984 - 1986 | unsexed/mixed | various gears | absolute number measured |
| Java Sea (Central Java), Indonesia | 1979 - 1979 | unsexed/mixed | trawls | absolute number measured |
| Leyte Gulf, Philippines | 1983 - 1987 | unsexed/mixed | various gears | absolute number measured |
| Manila Bay, Philippines | 1978 - 1979 | unsexed/mixed | trawls | absolute number measured |
| Samar Sea, Philippines | 1979 - 1979 | unsexed/mixed | trawls | absolute number measured |
| Southwest coast (Phuket, Phang-ga, Krabi, Trang & Satun), Thailand | 1984 - 1986 | unsexed/mixed | seines | raised to the catch |
| Tanjung Satai (southwest coast), Indonesia | 1971 - 1972 | unsexed/mixed | seines | % of sample |
| Visayan Sea, Philippines | 1983 - 1988 | unsexed/mixed | various gears | absolute number measured |

16) Morphometrics

Morphometric Data for *Rastrelliger brachysoma*

n = 1

| Picture Name | Length | | Lifestage | Aspect ratio |
|------------------------------|--------|----|-----------|--------------|
| Rabra_u0.jpg | 28.8 | SL | unsexed | 3.10 |
| | | | | |

17) Morphology

Morphometric Data for *Rastrelliger brachysoma*

| | |
|----------------------|--------------------------------------|
| Main Ref. | Collette, B.B., 2001 |
| Appearance refers to | |

| | |
|--------------------|--|
| Bones in OsteoBase | |
|--------------------|--|

Sex attributes

| | |
|----------------------|---------------------|
| Specialized organs | no special organs |
| Different appearance | males alike females |
| Different colors | males alike females |
| Remarks | |

Descriptive characteristics of juvenile and adult

| | |
|------------------------|--|
| Striking features | striking shape of body |
| Body shape lateral | fusiform / normal |
| Cross section | oval |
| Dorsal head profile | more or less straight |
| Type of eyes | more or less normal |
| Type of mouth/snout | more or less normal |
| Position of mouth | terminal |
| Type of scales | |
| Diagnosis | This species is distinguished by the following characters: body very deep, its depth at posterior margin of opercle 3.7-4.3 times in fork length; head equal to or less than body depth; maxilla covered by lacrimal bone but extending nearly to end of lacrimal; gill rakers very long, visible when mouth is opened, 30-48 on lower limb of first gill arch; numerous bristles on longest gill raker, about 150 on one side in specimens of 12.7 cm, 210 in specimens of 16 cm, and 240 at 19 cm fork length; intestine very long, 3.2-3.6 times fork length; snout pointed; swim bladder present; vertebrae 13 + 18 = 31; interpelvic process small and single; anal spine rudimentary. Colour of spinous dorsal fin yellowish with a black edge, pectoral and pelvic fins dusky, other fins yellowish (Ref. 168, 9684). |
| Ease of Identification | |

Meristic characteristics of *Rastrelliger brachysoma*

| | |
|-------------------------------|-------------------|
| Lateral Lines | 1 Interrupted: No |
| Scales on lateral line | |
| Pored lateral line scales | |
| Scales in lateral series | |
| Scale rows above lateral line | |
| Scale rows below lateral line | |

| | |
|--------------------------------|---------|
| Scales around caudal peduncle | |
| Barbels | 0 |
| Gill clefts (sharks/rays only) | absent |
| Gill rakers | |
| on lower limb | 30 - 48 |
| on upper limb | |
| total | |
| Vertebrae | |
| preanal | 13 - 13 |
| total | 31 - 31 |

Fins**Dorsal fin(s)**

| | |
|-----------------|------------------------|
| Attributes | no striking attributes |
| Fins number | 2 |
| Finlets No. | Dorsal 5 |
| | Ventral 5 |
| Spines total | 8 - 11 |
| Soft-rays total | 12 - 12 |
| Adipose fin | absent |

Caudal fin

| | |
|------------|-----------------------------|
| Attributes | forked; more or less normal |
|------------|-----------------------------|

Anal fin(s)

| | |
|-----------------|---------|
| Fins number | 1 |
| Spines total | 0 - 0 |
| Soft-rays total | 12 - 12 |

Paired fins

| | |
|----------|---|
| Pectoral | Attributes more or less normal Spines 0 Soft-rays 19 - 20 |
| Pelvics | Attributes more or less normal |
| | Position thoracic before origin of D1 |
| | Spines |
| | Soft-rays |

18) Larvae

(NA)

19) Recruitment

(NA)

20) Abundance

(NA)

References

1. Aglen, A., L. Føyn, O.R. Godø, S. Myklevoll and O.J. Østvedt, 1981. A survey of the marine fish resources of the north and west coast of Sumatra, August 1980. Reports on Surveys with the R/V 'Dr. Fridtjof Nansen', Institute of Marine Research. Bergen. 55 p.
2. al Sakaff, H. and M. Esseen, 1999. Occurrence and distribution of fish species off Yemen (Gulf of Aden and Arabian Sea). Naga ICLARM Q. 22(1):43-47.
3. Allen, G.R. and M.V. Erdmann, 2013. Coral reef fishes of Timor-Leste. p. 33-82. In G.R. Allen and M.V. Erdmann (eds). A rapid biological assessment of Timor-Leste. Conservation International.
4. Anonymous, 1985. Report of the Second Working Group Meeting on the Mackerels (Decapterus and Rastrelliger spp.) in the Malacca Strait, 4-9 October 1985, Colombo, Sri Lanka. Bay of Bengal Programme Document. 23 p.
5. Anonymous, 1996. Fish collection database of the University of British Columbia Fish Museum Fish Museum. University of British Columbia, Vancouver, Canada.
6. Anonymous, 1998. Fish collection database of the Bernice P. Bishop Museum (BPBM). Bishop Museum, 1525 Bernice Street, Honolulu, Hawai'i, 96817-0916 USA.
7. Anonymous, 1999. Fish collection database of the Natural History Museum, London (formerly British Museum of Natural History (BMNH)). Natural History Museum, London (formerly British Museum of Natural History (BMNH)).
8. Anonymous, 1999. Fish collection database of the Natural History Museum, London (formerly British Museum of Natural History (BMNH)). Natural History Museum, London (formerly British Museum of Natural History (BMNH)).
9. Anonymous, 2000. Fish collection database of the Gulf Coast Research Laboratory (GCRL). The Gulf Coast Research Laboratory (GCRL), Ocean Springs, Mississippi, USA.
10. Anonymous, 2000. The ichthyological collection of the Zoological Museum Hamburg (ZMH). Division of Ichthyology and Herpetology, Zoological Museum Hamburg (ZMH).
11. Anonymous, 2001. Fish collection database of the National Museum of Natural History (Smithsonian Institution), Smithsonian Institution-Division of Fishes.
12. Anonymous, 2002. Fish collection database of the American Museum of Natural History. American Museum of Natural History, Central Park West , NY 10024-5192, USA.
13. Anonymous, 2002. Fish collection database of the American Museum of Natural History. American Museum of Natural History, Central Park West , NY 10024-5192, USA.
14. Anonymous, 2002. Fish collection of the University of the Philippines in the Visayas Museum. UPV Museum.
15. Aprieto, V.L. and E.P. Viloso, 1979. Catch composition and relative abundance of trawl-caught fishes in the Visayan Sea. Fish. Res. J. Philipp. 4(1):9-18.

16. Aprieto, V.L. and E.P. Viloso, 1982. Demersal fish resources of Lingayen Gulf. Fish. Res. J. Phillips. 7(2):40-49.
17. Armada, N. and G. Silvestre, 1981. Demersal fish resource survey in Samar Sea and Carigara Bay. UP-NSDB Project 7811.1c Ag. 56 p. College of Fisheries, University of the Philippines in the Visayas, Iloilo City, Philippines.
18. Armada, N.B., 2004. Fish resource assessment and management recommendations for Davao Gulf. p. 332-335. In DA-BFAR (Department of Agriculture-Bureau of Fisheries and Aquatic Resources). In turbulent seas : the status of Philippine marine fisheries. CRMP document No: 02-CRM/2004. Cebu City, Philippines. 378 p.
19. Arthur, J.R. and S. Lumanlan-Mayo, 1997. Checklist of the parasites of fishes of the Philippines. FAO Fish. Tech. Pap. 369, 102 p. FAO, Rome.
20. Basmayor, L.O., R.D. Dioneda and V.S. Soliman, 1997. The fishes and invertebrates of San Miguel Bay. p.36-47. In V.S. Soliman and R.D. Dioneda (eds.) Capture Fisheries Assessment of San Miguel Bay, Post-Resource and Ecological Assessment of San Miguel Bay, Phil., Vol.1. BFAR, Fish. Sect. Prog. and Bicol Univ. Coll. of Fish. SMB Post-REA Tech. Rep. 1,60 p.
21. Beverton, R.J.H. and S.J. Holt, 1959. A review of the lifespans and mortality rates of fish in nature, and their relation to growth and other physiological characteristics. p. 142-180. In G.E.W. Wolstenholme and M. O'Connor (eds.) CIBA Foundation colloquia on ageing: the lifespan of animals. volume 5. J & A Churchill Ltd, London.
22. Bleeker, P., 1851. Over eenige nieuwe geslachten en soorten van Makreelachtige visschen van den Indischen Archipel. Natuurkd. Tijdschr. Neder.-Indië 1:341-372.
23. Boonraksa, V., 1987. Preliminary resource analysis of chub mackerel (*Rastrelliger* spp.) and round scads (*Decapterus* sp.) in the west coast of Thailand. Paper presented at the 3rd Working Group Meeting of the Malacca Strait Project/BOBP, 18-26 August 1986, Phuket, Thailand.
24. Boonraksa, V., 1988. Growth, mortality and maximum sustainable yield of the Indo-Pacific mackerel (*Rastrelliger brachysoma*) off the southwest coast of Thailand. FAO Fish. Rep. 389:356-371.
25. Broad, G., 2003. Fishes of the Philippines. Anvil Publishing, Inc., pasi City. 510 pp.
26. Cabanban, A.S., 1991. The dynamics of Leiognathidae in tropical demersal ichthyofaunal community. James Cook University of North Queensland, Australia. 262 p. Ph.D. dissertation.
27. Calud, A., E. Cinco and G. Silvestre, 1991. The gill net fishery of Lingayen Gulf, Philippines. p.45-50. In L.M.Chou, T.-E.Chua, H.W. Khoo, P.E. Lim, J.N.Paw, G.T.Silvestre, M.J.Valencia, A.T.White and P.K.Wong (eds.)Towards an integrated management of tropical coastal resources.ICLARM Conf.Proc.22, 455p.NUS, Sing.; NSTB, Sing.; and ICLARM, Phil.
28. Calumpang, H.P., L.J. Raymundo and E.P. Solis-Duran (eds.), 1994. Resource and ecological assessment of of Sogod Bay, Leyte, Philippines - Final Report Vol.1 Fisheries Assessment. Siliman University Marine Laboratory.
29. Calumpang, H.P., L.J. Raymundo, E.P. Solis-Duran and R.O. de Leon (eds.), 1993. Resource and ecological assessment of Carigara Bay, Leyte, Philippines - Final Report Vol.2 Fisheries resources and exploitation. Siliman University Marine Laboratory. 342 p.
30. Carl, H., 2003. Danish fish names. Zoological Museum of Copenhagen. Unpublished.
31. Cheung, W.L., R. Watson and D. Pauly, 2013. Signature of ocean warming in global fisheries catch. Nature 497:365-368.
32. Chinese Academy of Fishery Sciences, 2003. Chinese aquatic germplasm resources database. <http://zzzy.fishinfo.cn>

33. Chuenpagdee, R., 2002. Checklist of Thai names and scripts. Personal communication, April 2002.
34. Cinco, E., 1982. Length-weight relationships of fishes. p. 34-37. In D. Pauly and A.N. Mines (eds.) Small scale fisheries of San Miguel Bay, Philippines: biology and stock assessment. ICLARM Technical Reports 7, 124 p. Institute of Fisheries Development and Research, College of Fisheries, University of the Philippines in the Visayas, Quezon City, Philippines; International Center for Living Aquatic Resources Management, Manila, Philippines; and the United Nations University, Tokyo, Japan.
35. Cinco, E.A., 1996. Resource and ecological assessment of Sorsogon Bay, Philippines - Technical reports Vol.3 Capture fisheries assessment. Fisheries Sector Program Department of Agriculture-Asian Development Bank, United Business Technologies, Inc., Pasig Philippines. 158 p.
36. Cinco, E.A., J.C. Diaz, Q.P. Sia III and G.T. Silvestre, 1994. A checklist of fishes caught in San Miguel Bay, Philippines. In G. Silvestre, C. Luna and J. Padilla (eds.) Multidisciplinary assessment of the fisheries in San Miguel Bay, Philippines (1992-1993). ICLARM Technical Report 47. International Center for Living Aquatic Resources Management, Makati, Philippines.
37. Collette, B.B. and C.E. Nauen, 1983. FAO Species Catalogue. Vol. 2. Scombrids of the world. An annotated and illustrated catalogue of tunas, mackerels, bonitos and related species known to date. Rome: FAO. FAO Fish. Synop. 125(2):137 p.
38. Collette, B.B., 2001. Scombridae. Tunas (also, albacore, bonitos, mackerels, seerfishes, and wahoo). p. 3721-3756. In K.E. Carpenter and V. Niem (eds.) FAO species identification guide for fishery purposes. The living marine resources of the Western Central Pacific. Vol. 6. Bony fishes part 4 (Labridae to Latimeriidae), estuarine crocodiles. FAO, Rome.
39. Collette, B.B., 2003. Family Scombridae Rafinesque 1815 - mackerels, tunas, and bonitos. Calif. Acad. Sci. Annotated Checklists of Fishes (19):28.
40. Coppola, S.R., W. Fischer, L. Garibaldi, N. Scialabba and K.E. Carpenter, 1994. SPECIESDAB: Global species database for fishery purposes. User's manual. FAO Computerized Information Series (Fisheries). No. 9. Rome, FAO. 103 p.
41. Corpuz, A., J. Saeger and V. Sambilay, 1985. Population parameters of commercially important fishes in Philippine waters. Tech. Rep. Dep. Mar. Fish. Univ. Philipp. Visayas. (6):99 p.
42. Dalzell, P. and R.A. Ganaden, 1987. A review of the fisheries for small pelagic fishes in Philippine waters. Tech. Pap. Ser. Bur. Fish. Aquat. Resour. (Philipp.) 10(1):58 p. Bureau of Fisheries and Aquatic Resources, Quezon City, Philippines.
43. Dantis, A.L. and P.M. Aliño (comps.), 2002. Checklist of Philippine reef fishes. p. 208-226. In Aliño, P.M., E.F.B. Miclat, C.L. Nañola Jr., H.A. Roa-Quiaoit and R.T. Campos (eds.) Atlas of Philippine coral reefs. Philippine Coral Reef Information (Philreefs). Goodwill Trading Co., Inc. (Goodwill Bookstore), Quezon City, Philippines. xvi + 264p.
44. David, G., 1985. Pêche de subsistance et milieu naturel: les mangrove de Vanuatu et leur intérêt halieutique. Notes et documents d'océanographie. Mission ORSTOM de Port-Vila, 13:67 p. multigr.
45. Department of Fisheries Malaysia, 2009. Valid local name of Malaysian marine fishes. Department of Fisheries Malaysia. Ministry of Agriculture and Agro-based Industry. 180 p.
46. Department of Fisheries, 1987. Annual fishery statistics. Ministry of Agriculture, Malaysia.
47. Dioneda, R.R., L.R. Pura, Q.P. Sia III and L.O. Basmayor, 1995. A checklist of fishes and invertebrates caught and observed in Lagonoy Gulf. In G. Silvestre, C. Luna, V. Soliman

- and L. Garces (eds.) Resource and ecological assessment of Lagonoy Gulf, Philippines: Terminal Report Vol.2 Technical Monograph. ICLARM Tech Rep. 000,000 p.
48. Druzhinin, A.D. and D.T.T. Myint, 1970. A morphometric study of *Rastrelliger* spp. from the Mergui Archipelago. Proc. IPFC 13(2):49-58.
 49. Duong, T.T., 2001. Mot so loai ca thuong gap o bien Viet Nam (Viet Nam's Common Marine Fishes Catalogue). Ministry of Fisheries of Viet Nam-Fisheries Information Center of Viet Nam. 195 p.
 50. Dwiponggo, A., T. Hariati, S. Banon, M.L.D. Palomares and D. Pauly, 1986. Growth, mortality and recruitment of commercially important fishes and penaeid shrimps in Indonesian waters. ICLARM Tech. Rep. 17, 91 p.
 51. Eschmeyer, W.N. (ed.), 2001. Catalog of fishes. Updated database version of December 2001. Catalog databases as made available to FishBase in December 2001.
 52. Espejo-Hermes, J., 1998. Fish processing technology in the tropics. Tawid Publications, Quezon City, Philippines. 336 p.
 53. FAO, 1992. FAO yearbook. Fishery statistics: catches and landings. Vol. 70: 1990. FAO Fish. Ser. 38:647 p.
 54. FAO, 1994. FAO Yearbook. Fishery statistics: catches and landings. Vol. 74: 1992. FAO Fish. Ser. 43:677 p.
 55. FAO, 1995. FAO yearbook 1993. Fishery statistics. Catches and landings. FAO Fisheries Series No. 44. FAO Statistics Series No. 123. Vol. 76. 687 p.
 56. FAO-FIES, 2008. Aquatic Sciences and Fisheries Information System (ASFIS) species list. Retrieved from <http://www.fao.org/fishery/collection/asfis>, 29 April 2008.
 57. FAO-FIES, 2010. Aquatic Sciences and Fisheries Information System (ASFIS) species list. Retrieved from <http://www.fao.org/fishery/collection/asfis/en>, March 2010.
 58. FAO-FIES, 2012. Aquatic Sciences and Fisheries Information System (ASFIS) species list. Retrieved from <http://www.fao.org/fishery/collection/asfis/en>, March 2012.
 59. FAO-FIES, 2014. Aquatic Sciences and Fisheries Information System (ASFIS) species list. Retrieved from <http://www.fao.org/fishery/collection/asfis/en>, April 2014.
 60. FAO-FIES, 2015. Aquatic Sciences and Fisheries Information System (ASFIS) species list. Retrieved from <http://www.fao.org/fishery/collection/asfis/en>, [accessed 13/04/2015].
 61. FAO-FIES, 2017. Aquatic Sciences and Fisheries Information System (ASFIS) species list. Retrieved from <http://www.fao.org/fishery/collection/asfis/en> (accessed 08/06/2017).
 62. FAO-FIGIS, 2005. A world overview of species of interest to fisheries. Chapter: *Rastrelliger brachysoma*. Retrieved on 06 July 2005, from www.fao.org/figis/servlet/species?fid=2477. 2p. FIGIS Species Fact Sheets. Species Identification and Data Programme-SIDP, FAO-FIGIS
 63. Federizon, R.R., 1992. Description of the subareas of Ragay Gulf, Philippines, and their fish assemblages by exploratory data analysis. Aust. J. Mar. Freshwat. Res. 43:379-391.
 64. Ganaden, S.R. and F. Lavapie-Gonzales, 1999. Common and local names of marine fishes of the Philippines. Bureau of Fisheries and Aquatic Resources, Philippines. 385 p.
 65. Ganaden, S.R. and R.R. Calvelo, 1976. On the preliminary hydro-biological and fisheries survey of Estancia waters and its approaches in northern Iloilo. Phil. J. Fisheries. 13(2):163-177.
 66. Garces, L.R., M.L.D. Palomares and G.T. Silvestre, 1995. A preliminary trophic model of the coastal fisheries resources of Lagonoy Gulf. In G. Silvestre, C. Luna, V. Soliman and L. Garces (eds.) Resource and ecological assessment of Lagonoy Gulf, Philippines: Terminal Report Vol.2 Technical Monograph. ICLARM Tech Rep. 000,000 p.
 67. Gibbons, S., 1999. Collect fish on stamps. Stanley Gibbons Ltd., London and Ringwood. 418 p.

68. Gloerfelt-Tarp, T. and P.J. Kailola, 1984. Trawled fishes of southern Indonesia and northwestern Australia. Australian Development Assistance Bureau, Australia, Directorate General of Fishes, Indonesia, and German Agency for Technical Cooperation, Federal Republic of Germany. 407 p.
69. Goo, F.C.C. and A.H. Banner, 1963. A preliminary compilation of Fijian animal and plant names. Manuscript prepared under Contract SA-43-ph-3741. National Institute of Neurological Diseases and Blindness, National Institute of Health, Bethesda, Maryland, USA.
70. Grabda, E. and T. Heese, 1991. Polskie nazewnictwo popularne kraglouste i ryby. Cyclostomata et Pisces. Wyższa Szkoła Inżynierska w Koszalinie. Koszalin, Poland. 171 p. (in Polish).
71. Helfman, G.S. and J.E. Randall, 1973. Palauan fish names. Pac. Sci. 27(2):136-153.
72. Herre, A.W.C.T. and A.F. Umali, 1948. English and local common names of Philippine fishes. U. S. Dept. of Interior and Fish and Wildl. Serv. Circular No. 14, U. S. Gov't Printing Office, Washington. 128 p.
73. Herre, A.W.C.T., 1953. Check list of Philippine fishes. Res. Rep. U.S. Fish Wild. Serv. (20):1-977.
74. Hla Win, U., 1987. Checklist of fishes of Burma. Ministry of Livestock Breeding and Fisheries, Department of Fisheries, Burma.
75. Holt, S.J., 1959. Water temperature and cod growth-rate. J. Cons. Int. Explor. Mer. 24(3):374-376.
76. Hongskul, V., 1974. Population dynamics of pla-tu Rastrelliger neglectus (van Kampen) in the Gulf of Thailand. 15th Proc. IPFC, III:297-350.
77. Hureau, J.-C., 1991. La base de données GICIM : Gestion informatisée des collections ichthyologiques du Muséum. p. 225-227. In Atlas Préliminaire des Poissons d'Eau Douce de France. Conseil Supérieur de la Pêche, Ministère de l'Environnement, CEMAGREF et Muséum national d'Histoire naturelle, Paris.
78. Ingles, J. and D. Pauly, 1984. An atlas of the growth, mortality and recruitment of Philippines fishes. ICLARM Tech. Rep. 13. 127 p. International Center for Living Aquatic Resources Management, Manila, Philippines.
79. Isa, M.B.M., 1988. Population dynamics of *Nemipterus japonicus* (Pisces: Nemipteridae) off Kedah State, Malaysia. p. 126-140. In S.C. Venema, J.M. Christensen, and D. Pauly (eds.) Contributions to tropical fisheries biology. FAO/DANIDA Follow-up Training Course on Fish Stock Assessment in the Tropics, Denmark, 1986 and Philippines, 1987. FAOFish. Rep. (389).
80. Isa, M.M., 1987. On the status of the Rastrelliger and Decapterus fisheries of the west coast of peninsular Malaysia in 1984-1985. p. 81-100. In Investigations on the mackerel and scad resources of the Malacca Straits. Bay of Bengal Programme. BOBP/REP/39.
81. Kailola, P.J., 1991. The fishes of Papua New Guinea: a revised and annotated checklist. Vol. III. Gobiidae to Molidae. Research Bulletin No. 41, Research Section, Dept. of Fisheries and Marine Resources, Papua New Guinea. 153 p.
82. Kapoor, D., R. Dayal and A.G. Ponniah, 2002. Fish biodiversity of India. National Bureau of Fish Genetic Resources Lucknow, India. 775 p.
83. Kimura, S., 1995. A check list of the marine fishes collected around northern Palawan and Calauit islands, Philippines. p. 158-167. In Pawikan Conservation Project-PAWB, DENR, Philippines and Toba Aquarium, Japan. Dugongs Dugong dugon (Müller 1776) of the Philippines. A report of the Joint Dugong Research and Conservation Program. May 1995.
84. Klawe, W.L., 1980. Classification of the tunas, mackerels, billfishes, and related species, and their geographical distribution. Inter-Amer. Trop. Tuna Comm., Spec. Rep. 2:5-16.

85. Kurogane, K., 1974. Review of the mackerel resources of the western Gulf of Thailand. Proc. IPFC 15(3):253-264.
86. Kuronuma, K., 1961. A check list of fishes of Vietnam. United States Consultants, Inc.; International Cooperation Administration Contract-IV-153. Division of Agriculture and Natural Resources, United States Operations Mission to Vietnam. 66 p.
87. Lavapie-Gonzales, F., S.R. Ganaden and F.C. Gayanilo Jr., 1997. Some population parameters of commercially important fishes in the Philippines. Bureau of Fisheries and Aquatic Resources, Philippines. 114 p.
88. McManus, L.T. and T.-E. Chua (eds.), 1990. The coastal environmental profile of Lingayen Gulf, Philippines. ICLARM Technical Reports 22, 69 p. International Center for Living Aquatic Resources Management, Manila, Philippines.
89. Mindanao State University at Naawan, 1995. Rapid resource appraisal of Davao Gulf. Mindanao State University at Naawan Foundation for Science and Techology Development, Inc.
90. Mohsin, A.K.M. and M.A. Ambak, 1996. Marine fishes and fisheries of Malaysia and neighbouring countries. University of Pertanian Malaysia Press, Serdang, Malaysia. 744 p.
91. Mohsin, A.K.M., M.A. Ambak and M.N.A. Salam, 1993. Malay, English, and scientific names of the fishes of Malaysia. Occas. Publ. Fac. Fish. Mar. Sci. Univ. Pertanian Malays. 11:226 p.
92. Monkprasit, S., S. Sontirat, S. Vimollohakarn and T. Songsirikul, 1997. Checklist of Fishes in Thailand. Office of Environmental Policy and Planning, Bangkok, Thailand. 353 p.
93. Monkprasit, S., S. Sontirat, S. Vimollohakarn and T. Songsirikul, 1997. Checklist of Fishes in Thailand. Office of Environmental Policy and Planning, Bangkok, Thailand. 353 p.
94. Myers, R.F., 1999. Micronesian reef fishes: a comprehensive guide to the coral reef fishes of Micronesia, 3rd revised and expanded edition. Coral Graphics, Barrigada, Guam. 330 p.
95. Nomura, H. and M.S. de Sousa Rodriguez, 1967. Biological notes on king mackerel, *Scomberomorus cavalla* (Cuvier), from northeastern Brazil. Arq. Estac. Biol. Mar. Univ. Ceará, Fortaleza 7(1):79-85.
96. Ochavillo, D. and G. Silvestre, 1991. Optimum mesh size for the trawl fisheries of Lingayen Gulf, Philippines. p.41-44. In L.M.Chou,T.-E.Chua, H.W. Khoo, P.E. Lim, J.N.Paw, G.T.Silvestre, M.J.Valencia, A.T.White and P.K.Wong (eds.). Towards an integrated management of tropical coastal resources.ICLARM Conf.Proc.22, 455p.NUS, Sing.; NSTB, Sing.; and ICLARM, Phil.
97. Ochavillo, D., A. Calud and G. Silvestre, 1991. Population parameters and exploitation rates of trawl-caught fish species in Lingayen Gulf, Philippines. p.37-40. In L.M.Chou,T.-E.Chua,H.W. Khoo,P.E. Lim, J.N.Paw, G.T.Silvestre, M.J.Valencia, A.T.White and P.K.Wong (eds.). Towards an integrated management of tropical coastal resources.ICLARM Conf.Proc.22, 455p.NUS, Sing.; NSTB, Sing.; and ICLARM, Phil.
98. Ochavillo, D., H. Hernandez, S. Resma and G. Silvestre, 1989. Preliminary results of a study of the commercial trawl fisheries in Lingayen Gulf. p.31-42. In G. Silvestre, E. Milat and T.-E. Chua (eds.) Towards sustainable development of the coastal resources of Lingayen Gulf, Philippines. ICLARM Conference Proceedings 17, 200 p. PCAMRD, Los Baños, Lag. and ICLARM, Makati, Metro Manila,Phil.
99. Organisation for Economic Co-operation and Development, 1990. Multilingual dictionary of fish and fish products. Fishing News Books, Oxford.
100. Padilla, J.E., 1991. Managing tropical multispecies fisheries with multiple objectives. Ph.D. thesis, Simon Fraser University. 235 p.

101. Palomares, M.L.D., R. Froese, B. Derrick, S.-L. Nöel, G. Tsui, J. Woroniak and D. Pauly, 2018. A preliminary global assessment of the status of exploited marine fish and invertebrate populations. A report prepared by the Sea Around Us for OCEANA. The University of British Columbia, Vancouver, p. 64.
102. Pauly, D. and S. Aung, 1984. Population dynamics of some fishes of Burma based on length-frequency data. Bur/77/003/. FAO Field Doc. No. 7. 22 p. FAO, Rome.
103. Pauly, D., 1980. On the interrelationships between natural mortality, growth parameters, and mean environmental temperature in 175 fish stocks. *J. Cons. Int. Explor. Mer.* 39(2):175-192.
104. Pauly, D., 1982. The fishes and their ecology. p.15-33. In D. Pauly and A.N. Mines (eds.) Small-scale fisheries of San Miguel Bay, Philippines: biology and stock assessment. ICLARM Techn. Rep. 7, 124 p.
105. Pihu, E., 1979. Loomade elu 4. köide. Kalad Pihu, E. 1979. Loomade Elu, 4. Köide, Kalad. Valgus, Tallinn.
106. Pimoljinda, J., 1978. Study on the length-weight relationship of chub-mackerel (*Rastrelliger neglectus*) on the Indian coast of Thailand, 1967-1977. Annual Report 1978. Phuket Marine Fisheries Station, Phuket, Thailand. 11 p. (in Thai).
107. Pollnac, R.B. and M.L.G. Gorospe, 1998. Aspects of the human ecology of the coral reefs of Atulayan Bay. p. 77-139. In R.B. Pollnac (ed.) Rapid assessment of management parameters for coral reefs. Coastal Resources Center, University of Rhode Island. CMR 2205, ICLARM 1445.
108. Rajan, P.T., C.R. Sreeraj and T. Immanuel, 2011. Fish fauna of coral reef, mangrove, freshwater, offshore and seagrass beds of Andaman and Nicobar Islands. Zoological Survey of India, Andaman and Nicobar Regional Centre, Haddo, Port Blair.
109. Randall, J.E. and K.K.P. Lim (eds.), 2000. A checklist of the fishes of the South China Sea. *Raffles Bull. Zool. Suppl.* (8):569-667.
110. Rau, N. and A. Rau, 1980. Commercial marine fishes of the Central Philippines (bony fish). German Agency for Technical Cooperation, Germany. 623 pp.
111. Resource Combines and Manuel S. Enverga University Foundation, 2001. Resource and ecological assessment of Tayabas Bay - Final Report Vol.1 Introduction and assessment of fishery resources. Bureau of Fisheries and Aquatic Resources, Fisheries Sector Program, Department of Agriculture. 306 p.
112. Ricker, W.E., 1973. Russian-English dictionary for students of fisheries and aquatic biology. Fisheries Research Board of Canada, Ottawa.
113. Riede, K., 2004. Global register of migratory species - from global to regional scales. Final Report of the R&D-Projekt 808 05 081. Federal Agency for Nature Conservation, Bonn, Germany. 329 p.
114. Robins, C.R., R.M. Bailey, C.E. Bond, J.R. Brooker, E.A. Lachner, R.N. Lea and W.B. Scott, 1991. World fishes important to North Americans. Exclusive of species from the continental waters of the United States and Canada. *Am. Fish. Soc. Spec. Publ.* (21):243 p.
115. Sambilay, V.C. Jr., 1991. Depth-distribution patterns of demersal fishes of the Samar Sea, Philippines, and their use for estimation of mortality. M.Sc. thesis, University of the Philippines in the Visayas. 66 p.
116. Sanches, J.G., 1989. Nomenclatura Portuguesa de organismos aquáticos (proposta para normalização estatística). *Publ. Avuls. Inst. Nac. Invest. Pescas* 14:322 p.
117. Schuster, W.H. and R. Djajadiredja, 1952. Local common names of Indonesian fishes. W.V. Hoeve, Bandung, Indonesia. 276 p.

118. Sia, Q.P. III, L.R. Garces and E.A. Cinco, 1994. Reef fisheries of northwestern San Miguel Bay. In G. Silvestre, C. Luna and J. Padilla (eds.) Multidisciplinary assessment of the fisheries in San Miguel Bay, Philippines (1992-1993). ICLARM Technical Report 47. International Center for Living Aquatic Resources Management, Makati, Philippines.
119. Silvestre, G.T., L.R. Garces and C.Z. Luna, 1995. Resource and ecological assessment of Lagonoy Gulf, Philippines: Terminal Report Vol.1 Final Report. In G. Silvestre, C. Luna, V. Soliman and L. Garces (eds.) Resource and ecological assessment of Lagonoy Gulf, Philippines. ICLARM Tech. Rep. 000, 000 p.
120. Soliman, V.S., P. Nieves, L.R. Garces and Q.P. Sia III, 1995. Catch and effort in the Lagonoy Gulf fisheries. p. 000-000. In G. Silvestre, C. Luna, V. Soliman and L. Garces (eds.) Resource and ecological assessment of Lagonoy Gulf, Philippines, Vol. 2 (Technical Monograph). ICLARM Tech. Rep. 000, 000 p.
121. Somjaiwong, D. and S. Chullasorn, 1974. Tagging experiments on the Indo-Pacific mackerel *Rastrelliger neglectus* (van Kampen) in the Gulf of Thailand (1960-1965). Proc. Indo-Pac. Coun. 15(3):287-296.
122. Subramaniam, S., 1992. Fishery resources. Singapore Journal of Tropical Geography 13(1): 52-62.
123. Sucondharman, P., C. Tantisawetrat and U. Sriruangcheep, 1970. Estimation of age and growth of chub mackerel *Rastrelliger neglectus* (van Kampen) in the Western Gulf of Thailand. p. 471-480. In J.C. Marr (ed.) The Kuroshio: a symposium on the Japanese current. East West Center Press, Honolulu. 614 p.
124. Sudjastani, T., 1974. The species of *Rastrelliger* in the Java Sea, their taxonomy, morphometry and population dynamics. University of British Columbia. 147 p. Thesis.
125. Tandog-Edralin, D., S.R. Ganaden and P. Fox, 1988. A comparative study of fish mortality rates in moderately and heavily fished areas of the Philippines. p. 468-481. In S.C. Venema, J.M. Christensen and D. Pauly (eds.) Contributions to tropical fisheries biology. FAO/DANIDA Follow-up Training Course on Fish Stock Assessment in the Tropics, Denmark, 1986 and Philippines, 1987. FAO Fish. Rep. (389).
126. Thaman, R.R., T. Fong and A. Balawa, 2008. Ilava Ni Navakavu: Finfishes of Vanua Navakavu, Viti Levu, Fiji Islands. SPRH-FIO Biodiversity and Ethnobi多样性 Report No. 4, The University of the South Pacific, Suva, Fiji Islands.
127. Thomas, R.C., P.D. Beldia III and W.L. Campos, 2013. Resolving the identity of larval fishes, Dulong, in the Verde Island Passages, Philippines. Phil. J. Nat. Sci. 18:29-36.
128. Torres, F. Jr., 2017. Species list for the five municipalities in Southern Palawan. Personal Communication [31/03/17].
129. U.P. Visayas Foundation, 1993. Resource and ecological assessment of Ormoc Bay. Vol.5. The fisheries of Ormoc Bay. Institute of Marine Fisheries and Oceanology Bureau of Fisheries and Aquatic Resources. IMFO Tech. Rep. No.15.
130. University of the Philippines at Los Baños Foundation, 1995. Resource and ecological assessment of Ragay Gulf Vol.3 Summary tables and figures. University of the Philippines at Los Baños Foundation, Inc.
131. University of the Philippines at Los Baños Foundation, 1996. Resource and ecological assessment of Ragay Gulf Vol.4 Raw data tables. University of the Philippines at Los Baños Foundation, Inc.
132. Viloso, E.P., G.V. Hermosa and C. Dizon, 1983. Species composition and diversity of fishes caught by otter trawling in Samar Sea. Fish. Res. J. Philipp. 8(2):33-49.
133. Wass, R.C., 1984. An annotated checklist of the fishes of Samoa. Natl. Ocean. Atmos. Adminis. Tech. Rept., Natl. Mar. Fish. Serv., Spec. Sci. Rept. Fish. (781).

134. Wu, H.L., K.-T. Shao and C.F. Lai (eds.), 1999. Latin-Chinese dictionary of fishes names. The Sueichan Press, Taiwan. 1028 p.
135. Ziegler, B., 1979. Growth and mortality rates of some fishes of Manila bay. Philippines as estimated from the analysis of length frequencies. Mathematisch-Naturwissenschaftliche Fakultät der Christian-Albrechts-Universität, Kiel 117p. Thesis.
136. Collette, B.; Di Natale, A.; Fox, W.; Juan Jorda, M. & Nelson, R. (2011). "*Rastrelliger brachysoma*". IUCN Red List of Threatened Species. IUCN. 2011: e.T170318A6745895. doi:10.2305/IUCN.UK.2011-2.RLTS.T170318A6745895.en.
137. ^ "*Rastrelliger brachysoma* (Bleeker, 1851)". Integrated Taxonomic Information System. Retrieved 5 August 2011.
138. ^ Jump up to:^{a b} Froese, Rainer and Pauly, Daniel, eds. (2015). "*Rastrelliger brachysoma*" in FishBase. 12 2015 version.
139. ^ Jump up to:^{a b} Sukanan, Darunee (18 July 2019). "Illegal fishing endangers both ecosystems and food security". Sustainability Times. Retrieved 19 July 2019.
140. ^ "*Holy Mackerel*". Archived from the original on 2011-11-20. Retrieved 2011-11-20.
141. ^ Pla thu is a very Thai fish Archived 2013-01-22 at the Wayback Machine
142. ^ WONGCHA-UM, PANU (2010). WHAT IS THAI CUISINE? THAI CULINARY IDENTITY CONSTRUCTION FROM THE RISE OF THE BANGKOK DYNASTY TO ITS REVIVAL (MA Thesis). History Department, National University of Singapore. Retrieved 27 August 2017. [dead link]
143. ^ Akkasit Jongjareonrak et al. Antioxidant activity of fermented fish viscera (Tai-Pla) from short-bodied mackerel, Faculty of Agro-Industry, Prince of Songkla University, Hat Yai, Songkhla, 90112 Thailand
144. ^ Kongrut, Anchalee (21 December 2019). "Humble 'pla tu' caught up in nets of destruction" (Opinion). Bangkok Post. Retrieved 21 December 2019.